


Sammanfattning

Vårens kiselalgsblomning hade startat vid den södra Kattegattstationen Anholt E. Stormen Egon hindrade provtagning i Skagerrak under denna expedition och flertalet stationer i Östersjön fick strykas av samma orsak.

I Östersjön var både artdiversiteten och celltätheten mycket låg vid alla besökta stationer.

De integrerade (0-10 m) klorofyll α värdena låg inom det normala för denna månad i de besökta havsområdena.

Fluorometern på ctdn var på service så inga fluorescensdata är med i klorofylldiagrammen på s 6 - 9.



Abstract

The diatom spring bloom had started at the southern Kattegat station, Anholt E. Due to the storm Egon, the January cruise could not visit the Skagerrak area this month, as well as several stations in the Baltic.

The species diversities and cell densities were very low in the Baltic Sea.

The integrated (0-10 m) chlorophyll α concentrations were normal for this month.

The ctd fluorometer was being serviced why there are no fluorescence data in the chlorophyll diagrams on p. 6 - 9.

More detailed information on species composition and abundance

The Skagerrak

Å17 (open Skagerrak) and Släggö (Skagerrak coast)

Due to the storm Egon the area was not visited at all during the expedition.

The Kattegat

Anholt E 12th of January

The diatom spring bloom had started at Anholt E. Diatoms dominated the phytoplankton community, *Skeletonema marinoi* and the potentially toxic genus *Pseudo-nitzschia* spp were very common and numerous.

The integrated (0-10 m) chlorophyll α concentrations were normal in the Kattegat for this month.




Fig.1 The spring bloom had already started in the southern part of the Kattegat area at the time of the January cruise.




Fig.2. The beautiful phytobenthic diatom cf. *Gyrosigma* sp. visited the pelagic part of the station Anholt E.




Fig.3 The spring bloom had not started in the northern part of the Kattegat and the diatom species *Leptocylindrus danicus* was rather alone in the sample from N14.

The Baltic Sea

The species diversity and cell densities were very low in the Baltic Sea. There were only a few cells of each of the species registered in Table 2.

The integrated (0-10 m) chlorophyll α concentrations were normal in the Baltic Sea for this month.





Fig.4 Very few cells were found in the Baltic Sea. This cyanobacterial colony, cf. *Aphanocapsa* sp., was found in BCSIII, the South East Baltic Sea.


Selection of observed species	Anholt E	N14 Falkenberg
Red=potentially toxic species	12/1	12/1
Hose 0-10 m	presence	presence
<i>Cerataulina pelagica</i>		present
<i>Chaetoceros danicus</i>		present
<i>Coscinodiscus centralis</i>	present	
<i>Coscinodiscus radiatus</i>	present	
<i>Ditylum brightwellii</i>	common	
<i>Eucampia zodiacus</i>	common	
<i>Guinardia delicatula</i>	common	
<i>Nitzschia longissima</i>		present
<i>Proboscia alata</i>	common	present
Pseudo-nitzschia spp	very common	present
<i>Pseudosolenia calcar-avis</i>	present	
<i>Skeletonema marinoi</i>	very common	common
<i>Thalassiosira spp</i>	common	present
<i>Thalassiosira angulata</i>	common	common
<i>Thalassiosira anguste-lineata</i>	common	
<i>Thalassiosira punctigera</i>	present	
<i>Ceratium furca</i>	common	
<i>Ceratium fusus</i>	common	present
<i>Ceratium lineatum</i>	common	present
<i>Ceratium longipes</i>	common	
<i>Ceratium tripos</i>	common	present
Dinophysis norvegica	present	
<i>Gyrodinium spp</i>	present	
<i>Polykrikos schwartzii</i>	present	
<i>Protoperidinium spp</i>	present	
<i>Protoperidinium bipes</i>	present	
<i>Dictyocha fibula</i>	common	
Dictyocha speculum	common	
<i>Woronichinia spp</i>		present

Selection of observed species	BCS III-10	BY2	BY15	BY31	BY38
Red=potentially toxic species	13/1	13/1	14/1	8/1	9/1
Hose 0-10 m	presence	presence	presence	presence	presence
Chaetoceros spp					present
Chaetoceros danicus	present	present	present	present	
Nitzschia longissima		present			
Skeletonema marinoi		present			present
Dinophysis acuminata			present	present	present
Dinophysis norvegica				present	present
Gymnodiniales			present	present	present
Gyrodinium spp			present		
Protoperidinium spp				present	
Aphanocapsa spp	present				
Woronichinia spp				present	present
Oocystis spp				present	
Pachysphaera spp			present		
Planctonema lauterbornii				present	present
Craspedophyceae				present	
Ebria tripartita				present	present
Cryptomonadales			present	present	present
Ciliophora			present		


The Kattegat and The Sound




The Southern Baltic



The Eastern Baltic



The Western Baltic



Om klorofylldiagrammen

Klorofyll a är ett mått på mängden växtplankton. Prover tas från ett antal djup. Data presenteras både från de fasta djuren och som medelvärdet 0-20 m. Utöver resultaten från laboratorieanalyserna av vattenprover mäts klorofyll a som fluorescens från ett automatiskt instrument som sänks ned från fartyget. På så sätt kan djupt liggande, ibland tunna lager av växtplankton observeras.

About the chlorophyll graphs

Chlorophyll a is sampled from several depths. Data are presented both from the discrete depths and as an average 0-20 m. In addition to the laboratory analysis from the water samples chlorophyll fluorescence is measured in continuous depth profiles from the ship. This is a way to observe thin layers of phytoplankton occurring below the surface.

Om AlgAware

SMHI genomför månatliga expeditioner i Östersjön och Västerhavet. Resultat baserade på semikvantitativ mikroskopanalys av planktonprover samt klorofyllmätningar presenteras kortfattat i denna rapport. Information från SMHIs satellitövervakning av algbloomingar finns under perioden juni-augusti på www.smhi.se.

About AlgAware

SMHI carries out monthly cruises in the Baltic and the Kattegat/Skagerrak. Results from semi quantitative microscopic analysis of phytoplankton samples as well as chlorophyll measurements are presented in brief in this report. Information from SMHIs satellite monitoring of algal blooms is found on www.smhi.se during the period June-August.

Art / Species	Gift / Toxin	Eventuella symptom	Clinical symptoms
<i>Alexandrium</i> spp.	Paralytic shellfish poisoning (PSP)	Milda symptom: Inom 30 min.: Stickningar eller en känsa av bedövning runt läpparna, som sprids gradvis till ansiktet och nacken; stickningar i fingertoppar och tår; Huvudvärk; yrsel, illamående, kräkningar, diarré Extrema symptom: Muskelförlamning; andningssvårigheter; känsa av att kvävas; Man kan vara död inom 2-24 timmar efter att ha fått i sig giften, på grund av att andningsmuskulaturen förlamas.	Mild case: Within 30 min: tingling sensation or numbness around lips, gradually spreading to face and neck; prickly sensation in fingertips and toes; headache, dizziness, nausea, vomiting, diarrhoea. Extreme case: Muscular paralysis; pronounced respiratory difficulty; choking sensation; death through respiratory paralysis may occur within 2-24 hours after ingestion.
<i>Dinophysis</i> spp.	Diarrehetic shellfish poisoning (DSP)	Milda symptom: Efter cirka 30 minuter till några timmar: yrsel, illamående, kräkningar, diarré, magont Extrema symptom: Upprepad exponering kan orsaka cancer	Mild case: Within 30 min-a few hours: dizziness, nausea, vomiting, diarrhoea, abdominal pain. Extreme case: Repeated exposure may cause cancer.
<i>Pseudo-nitzschia</i> spp.	Amnesic shellfish poisoning (ASP)	Milda symptom: Efter 3-5 timmar: yrsel, illamående, kräkningar, diarré, magkramper Extrema symptom: Yrsel, hallucinationer, förvirring, förlust av korttidsmindet, kramper	Mild case: Within 3-5 hours: dizziness, nausea, vomiting, diarrhoea, abdominal cramps. Extreme case: dizziness, hallucinations, confusion, loss of memory, cramps.
<i>Chaetoceros concavicornis/ C.convolutus</i>	Mechanical damage through hooks on setae	Låg celltäthet: Ingen påverkan. Hög celltäthet: Fiskens gälar skadas, fisken dör.	Low cell numbers: No effect on fish. High cell numbers: Fish death due to gill damage.
<i>Pseudochattonella</i> spp.	Fish toxin	Låg celltäthet: Ingen påverkan. Hög celltäthet: Fiskens gälar skadas, fisken dör.	Low cell numbers: No effect on fish. High cell numbers: Fish death due to gill damage.

Översikt över några potentiellt skadliga alger och det aktuella giftets effekt. Overview of potentially harmful algae and effects of toxins. Manual on harmful marine microalgae (2003 - UNESCO Publishing).

Kartan på framsidan visar viktat medelvärde för klorofyll α , $\mu\text{g/l}$ (0-20 m) vid de olika stationerna. Förekomst av skadliga alger vid stationer där arter analyseras markeras med symbol.

The map on the front page shows weighted mean of chlorophyll α , $\mu\text{g/l}$ (0-20 m) at sampling stations. Presence of harmful algae at stations where species analysis is performed is shown with a symbol.

