

Report from the SMHI monitoring cruise with R/V Aranda



Survey period: 2016-10-10 to 2016-10-17

Principal: Swedish Meteorological and Hydrological Institute, the Swedish Agency for Marine and Water Management and Finnish Environment Institute

SUMMARY

The cruise, which is part of the Swedish marine monitoring programme, covered the Skagerrak, the Kattegat, the Sound, the Gulf of Finland and the Baltic Proper.

The temperature in the surface water was now normal and the cooling of the surface water had started. This was clearly seen in the Skagerrak and Kattegat. The salinity showed normal values in the whole investigated area.

Generally, the nutrient concentrations in the surface water had increased since the last cruise in September, which is normal for the season. Though, high concentrations of inorganic nitrogen and phosphate were found in certain parts of the Baltic Proper. The silicate concentration in the whole Baltic Proper continued to be highly elevated.

Acute hypoxia ($O_2 < 2$ ml/l) was found at some stations in the Kattegat and the Sound already from 20 meters depth. In the Baltic Proper and the Gulf of Finland acute hypoxia was found from depth exceeding 60-70 meters depth. Anoxia ($O_2 = 0$ ml/l) was found in the Western and Northern Gotland basin and outer Gulf of Finland from depth over 80-90 meters, in the Eastern Gotland basin at intermediate depth and at the bottom. Due to a longer period with strong northerly/easterly winds and high air pressure over the Baltic, the water level is now low in the whole Baltic Sea. This provides favourable conditions for a new large inflow to the Baltic Sea, which is needed to improve the oxygen situation in the Baltic Proper.

Next regular cruise is scheduled to start 14th of November.

RESULTS

The cruise was conducted aboard the Finnish research vessel Aranda and started in Helsinki on the 10th of October and ended in the same port on 17th of October. The winds during the cruise were fresh to strong, mainly from northeast to east. Due to the strong winds and rough seas the stations Hanöbukten and REF M1V1 could not be visited.

During the expedition extra zooplankton samples were taken at the stations; BY15, BY5, BY2, Anholt E and Släggö, on behalf of Umeå University for future analysis of mercury.

At N Midsjöbanken two hydrophones were deployed for the Swedish Defence Research Agency. The instruments will measure marine noise and register “click” sounds from harbour porpoise.

This report is based on data that have passed a first quality control. When data are published at the SMHI data centre some values might have changed after further quality controls have been performed. Preliminary data from this cruise are published as soon as possible on the data centres webpage, normally within one or two weeks after a cruise.

Download data here: <http://www.smhi.se/klimatdata/oceanografi/havsmiljodata>

The Skagerrak

The temperature in the surface water was normal for the season and varied around 11-12°C. The cooling of the surface water had started and a shallow thermocline was found between 5-15 meters depth, while another thermocline was found deeper at 40-60 meters depth. The surface salinity was lower than normal in the outer and southern parts, while normal in the remaining areas. The lowest salinities were found at the station P2, ~21 psu and highest at Å15, ~33 psu. The halocline coincided with the upper thermocline and was weakly developed at the deeper thermocline.

All nutrients in the surface layer showed increasing concentrations compared to the previous cruise in September. However, the concentrations were lower than normal in the same areas as low salinities were found. The concentration of phosphate varied between 0.04-0.09 µmol/l and inorganic nitrogen (nitrite, nitrate and ammonia) had increased and varied around 0.32-1.71 µmol/l. The silicate concentrations showed values between 0.3 and 2.2 µmol/l.

The fluorescence from the CTD was low at all stations but some phytoplankton activated could be seen in the surface layer.

The bottom water was well oxygenated, even at the coastal station; Släggö.

The Kattegat and the Sound

The temperature of the surface water was now normal again and around 12°C. The cooling of the surface water had started also here and a thermocline at 10-20 meters depth separated the cooler surface water from the somewhat warmer water at intermediate depths. The surface salinity was generally lower than normal, due to a long period characterized by strong outflow from the Baltic Sea, and varied between 17 and 22 psu. In the Sound the salinity was around 9 psu. The halocline and thermocline coincided.

Similar to the conditions in Skagerrak, all nutrients had increased since the visit in September and the concentrations were overall normal for the season. The phosphate concentration was 0.1-0.4 $\mu\text{mol/l}$, silicate 2-12 $\mu\text{mol/l}$ and the inorganic nitrogen varied between 0.3 and 0.8 $\mu\text{mol/l}$. The concentrations were highest in the Sound. The nutrient concentrations increased below the halocline and showed generally higher values than normal in the deep water.

The oxygen concentrations in the deep water were lower than normal in the whole Kattegat. Acute hypoxia ($\text{O}_2 < 2 \text{ ml/l}$) was found in the Sound and at the coastal station; N14 already from 20 meters depth. Hypoxia was also found at the bottom at Anholt E.

The phytoplankton activity was low in the whole investigated area.

The Baltic Proper

The water temperature had decreased since the previous cruise in September and was now normal in the whole Baltic Proper. Surface temperatures were 9-14 °C, lowest in the Gulf of Finland and highest in the Arkona Basin. The salinity was just below normal in the Eastern Gotland Basin and normal or just over normal in the remaining areas. The lowest salinity was found in the Gulf of Finland, ~5 psu and highest in the Arkona Basin, ~8 psu. The halocline and the thermocline followed each other at about 60-70 meters depth in the Eastern Gotland Basin and in the Gulf of Finland. In other areas the stratification was found at 30-40 meters depth.

The concentrations of inorganic nitrogen was just over normal in the southeast, south and western Baltic Proper and varied between 0.7 and 1.0 $\mu\text{mol/l}$. In the Eastern Gotland Basin lower concentrations were found, around 0.4 $\mu\text{mol/l}$. The phosphate concentration was normal in the whole area, except from higher than normal concentrations in the Western Gotland Basin. The concentration of phosphate varied between 0.2-0.4 $\mu\text{mol/l}$. Silicate continued to be higher than normal in the whole Baltic Proper and concentrations ranged between 11 and 16 $\mu\text{mol/l}$. Below the halocline the concentrations of nutrients increase in the whole area, while lower concentration than normal of phosphate and inorganic nitrogen was found in the deeper parts of the Eastern Gotland Basin, probably due to the influence of oxygenated water that was found at those depths.

In the Western Gotland Basin the oxygen situation was still severe. Anoxic conditions were found already at depth exceeding 80-90 meters. In the Eastern Gotland Basin anoxic conditions was found at intermediate depth and at the bottom while anoxia was found from 80 meters in the Northern Gotland Basin and the outer parts of the Gulf of Finland. However, the deep water in the central parts of the Gulf of Finland, at LL7 was still oxygenated. Acute hypoxia was found from 60-80 meters depth in the whole Baltic Proper.

Due to a longer period of strong easterly and northerly winds and high air pressure over the Baltic Sea the water level in the whole area is low. This provides favourable conditions for a new large inflow to the Baltic Sea this winter, which is needed to improve the oxygen situation in the Baltic Proper.

The fluorescence measurements from the CTD showed low phytoplankton activity in the whole investigated area.





Figure 2. Transect showing oxygen and salinity through the Baltic Proper, from the Sound to the Eastern Gotland Basin.

PARTICIPANTS

Name		Institute
Martin Hansson	Chief Scientist	SMHI
Örjan Bäck		SMHI
Jenny Lycken		SMHI
Johanna Linders		SMHI
Daniel Simonsson		SMHI
Emilia Lalander		Swedish Defence Research Agency (FOI)

APPENDICES

- Track chart
- Table over stations, analysed parameters and number of sampling depths
- Map showing bottom oxygen concentrations
- Vertical profiles for normal monitoring stations
- Monthly average surface water plots for normal monitoring stations



Year: 2016

Ser no	Cru no	Stat code	Proj	Stat name	Lat	Lon	Start date yyyymmdd	Start time hhmm	Bottom depth m	Secchi depth m	Wind di ve	Air temp C	Air pres hPa	WCWI No elac de	T e e m p t	T a a l l	S s s s	P d d x	D h o o	H 2 h t	P t t t	N n n n	N n n n	A n a r o o k	A n a z n t y	S H C D P T P L Y A U C I e l r n o	S H C D P T P L Y A U C I e l r n o			
															p p	t t	t t	y y	s s	t t	i a	r r	r r	r r	k o k	m h c c c n g l	n l	a c n	e d m	
															- - - -	- -	b c b c	b c							- s	i o	_ t			
															t t t t	t t									s	n w	2 _	5 v		
															l d l d	l d									i			i		
																												s		
																												i		
0544	18	GFX29	SYK	LL7	5950.79	02450.27	2016-10-10	11:30	104		8 15.0	9	1033	2750 13	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0545	18	BPX60	SYK	LL12	5929.044	02253.71	2016-10-10	17:55	84		8 16.1	9	1032	9999 11	x x	- x	- -	x x	x x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0546	18	BPX62	SYK	LL15	5911.001	02144.81	2016-10-10	22:20	132		9 9.0	6	1033	9999 14	x x	- x	- -	x x	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	
0547	18	BPX35	SYK	BY29/LL19	5852.909	02019.672	2016-10-11	02:30	174		8 8.0	6	1032	9990 16	- x	- x	- x	x x	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	
0548	18	BPEX26	BAS	BY20 FÄRÖDJUPET	5759.89	01952.73	2016-10-11	07:35	196	7	8 9.6	6	1031	2740 17	x x	- x	- x	x x	x x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0549	18	BPEX21	BAS	BY15 GOTLANDSDJ	5718.729	02004.569	2016-10-11	12:20	239	7	6 9.3	7	1029	1140 19	x x	- x	x x	x x	x x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0550	18	BPEX13	BAS	BY10	5638.021	01935.089	2016-10-11	17:50	145		10 9.5	7	1029	9990 15	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0551	18	BPSE11	BAS	BCS III-10	5533.319	01824.012	2016-10-12	01:50	90		7 8.5	7	1026	9990 12	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0552	18	BPSB07	BAS	BY5 BORNHOLMSDJ	5515	01559.05	2016-10-12	09:00	91		9 9.5	9	1026	2850 12	x x	- x	x x	x x	x x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0553	18	BPSB06	BAS	BY4 CHRISTIANSÖ	5522.98	01520.033	2016-10-12	13:10	92		9 13.0	10	1027	2850 12	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0554	18	BPSA03	BAS	BY2 ARKONA	5458.27	01405.931	2016-10-12	19:10	47		9 8.9	11	1026	9999 8	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0555	18	BPSA02	BAS	BY1	5500.949	01318.05	2016-10-12	22:40	46		11 8.8	10	1026	9990 8	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0556	18	SOCX39	BAS	W LANDSKRONA	5552	01244.918	2016-10-13	04:15	53		8 7.4	7	1028	9990 9	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0557	18	KAEX29	BAS	ANHOLT E	5640.121	01206.668	2016-10-13	09:30	63	7	9 11.3	9	1029	1440 10	x x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0558	18	KANX50	BAS	N14 FALKENBERG	5656.402	01212.71	2016-10-13	12:00	32	7	9 9.4	9	1029	1230 7	x x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0559	18	KANX25	BAS	FLADEN	5711.556	01139.477	2016-10-13	15:00	83		9 10.4	9	1028	2730 12	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0560	18	SKEX18	BAS	Å17	5817.062	01030.258	2016-10-13	22:00	350		7 9.3	7	1030	9999 15	x x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0561	18	SKEX17	BAS	Å16	5816.029	01043.459	2016-10-13	23:55	203		8 8.9	6	1030	9999 13	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	x	
0562	18	SKEX16	BAS	Å15	5817.62	01050.709	2016-10-14	01:15	137		7 6.9	6	1030	9999 12	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0563	18	SKEX15	BAS	Å14	5818.932	01056.54	2016-10-14	02:20	111		8 7.4	6	1029	9999 11	- x	- x	- -	x -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	
0564	18	SKEX14	BAS	Å13	5820.369	01101.665	2016-10-14	03:20	107		10 7.4	6	1029	9999 10	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0565	18	FIBG27	BAS	SLÄGGÖ	5815.592	01126.14	2016-10-14	05:15	78		8 8.3	6	1028	2820 9	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0566	18	SKEX23	BAS	P2	5752.602	01117.519	2016-10-14	12:00	95	7	7 9.8	8	1025	1330 10	x x	- x	- x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0567	18	KAEX29	BAS	ANHOLT E	5640.12	01206.67	2016-10-14	19:50	63		10 16.5	7	1022	9999 10	x x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x					
0568	18	BPSE00	EXT	N MIDSJÖBANKEN	5612.998	01716.141	2016-10-15	01:50	23		13 12.8	7	1028	9990 6	- x	- x	- -	x -	- -	- -	- -	- -	- -	- -	- -	- -	- -	- -	x	
0569	18	BPWX45	BAS	BY38 KARLSÖDJ	5707.032	01740.118	2016-10-16	06:40	111		14 12.7	7	1031	2840 14	x x	- x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x				
0570	18	BPWX38	BAS	BY32 NORRKÖPING	5801.01	01759.069	2016-10-16	12:40	201	7	15 8.1	6	1034	2840 17	x x	- x	- x	x x	x x	- x	- x	x x	x x	- -	x	- - - - - - - - - - - x				


Bottom water oxygen concentration (ml/l)

Country: Finland

Ship: Aranda

Date: 20161010-20161016

Series: 0544-0570



STATION P2 SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 80 m)


O₂ ml/l

O₂ ml/l



Vertical profiles P2 October







— Mean 2001-2015 St.Dev. • 2016-10-14





STATION A13 SURFACE WATER (0-10m)

Annual Cycles

— Mean 2001-2015 St.Dev. • 2016




OXYGEN IN BOTTOM WATER (depth >= 75 m)



Vertical profiles A13 October

— Mean 2001-2015 St.Dev. • 2016-10-14




STATION A15 SURFACE WATER (0-10m)


Annual Cycles

— Mean 2001-2015 St.Dev. • 2016


Temperature °C




Salinity psu




PO₄ µmol/l




DIN µmol/l



SiO₃ µmol/l




O₂ saturation %




OXYGEN IN BOTTOM WATER (depth >= 125 m)

O₂ ml/l




O₂ ml/l



Vertical profiles A15 October

— Mean 2001-2015 St.Dev. • 2016-10-14




STATION A17 SURFACE WATER (0-10m)


Annual Cycles

— Mean 2001-2015 St.Dev. • 2016


Temperature °C




Salinity psu




PO₄ µmol/l




DIN µmol/l



SiO₃ µmol/l




O₂ saturation %




OXYGEN IN BOTTOM WATER (depth >= 300 m)

O₂ ml/l




O₂ ml/l



Vertical profiles A17 October

— Mean 2001-2015 St.Dev. • 2016-10-13



STATION FLADEN SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 65 m)


O₂ ml/l

O₂ ml/l



Vertical profiles FLADEN October

— Mean 2001-2015 St.Dev. • 2016-10-13



STATION ANHOLT E SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 45 m)


O₂ ml/l

O₂ ml/l



Vertical profiles ANHOLT E October

— Mean 2001-2015 St.Dev. • 2016-10-14



STATION W LANDSKRONA SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 40 m)


O₂ ml/l

O₂ ml/l



Vertical profiles W LANDSKRONA October







— Mean 2001-2015 St.Dev. • 2016-10-13





STATION BY1 SURFACE WATER (0-10m)

Annual Cycles

— Mean 2001-2015 St.Dev. • 2016




OXYGEN IN BOTTOM WATER (depth >= 40 m)



Vertical profiles BY1 October

— Mean 2001-2015 St.Dev. • 2016-10-12



STATION BY2 ARKONA SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 40 m)


O₂ ml/l

O₂ ml/l



Vertical profiles BY2 ARKONA October







— Mean 2001-2015 St.Dev. • 2016-10-12





STATION BY4 CHRISTIANSÖ SURFACE WATER (0-10m)

Annual Cycles

— Mean 2001-2015 St.Dev. • 2016




OXYGEN IN BOTTOM WATER (depth >= 80 m)



Vertical profiles BY4 CHRISTIANSÖ October

— Mean 2001-2015 St.Dev. • 2016-10-12



STATION BY5 BORNHOLMSDJ SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 80 m)


O₂ ml/l

O₂ ml/l



Vertical profiles BY5 BORNHOLMSDJ October


— Mean 2001-2015 St.Dev. • 2016-10-12




STATION BCS III-10 SURFACE WATER (0-10m)

Annual Cycles


— Mean 2001-2015
Temperature °C




— St.Dev. ● 2016
Salinity psu




PO₄ µmol/l




DIN µmol/l



SiO₃ µmol/l




O₂ saturation %




OXYGEN IN BOTTOM WATER (depth >= 80 m)

O₂ ml/l




O₂ ml/l



Vertical profiles BCS III-10 October

— Mean 2001-2015 ■ St.Dev. ● 2016-10-12



STATION BY10 SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 125 m)


O₂ ml/l

O₂ ml/l



Vertical profiles BY10 October

— Mean 2001-2015 ■ St.Dev. ● 2016-10-11



STATION BY15 GOTLANDSDJ SURFACE WATER (0-10m)


Annual Cycles

— Mean 2001-2015


■ St.Dev.

● 2016


Temperature °C




Salinity psu




PO₄ µmol/l




DIN µmol/l



SiO₃ µmol/l




O₂ saturation %




OXYGEN IN BOTTOM WATER (depth >= 225 m)

O₂ ml/l




O₂ ml/l



Vertical profiles BY15 GOTLANDSDJ October


— Mean 2001-2015 ■ St.Dev. ● 2016-10-11




STATION BY20 FÅRÖDJ SURFACE WATER (0-10m)

Annual Cycles


— Mean 2001-2015
Temperature °C




St.Dev. ● 2016
Salinity psu




PO₄ µmol/l




DIN µmol/l



SiO₃ µmol/l




O₂ saturation %




OXYGEN IN BOTTOM WATER (depth >= 175 m)

O₂ ml/l




O₂ ml/l



Vertical profiles BY20 FÄRÖDJ October

— Mean 2001-2015 St.Dev. ● 2016-10-11



STATION BY32 NORRKÖPINGSDJ SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 175 m)


O₂ ml/l

O₂ ml/l



Vertical profiles BY32 NORRKÖPINGSDJ October

— Mean 2001-2015 St.Dev. ● 2016-10-16



STATION BY38 KARLSÖDJ SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 100 m)


O₂ ml/l

O₂ ml/l



Vertical profiles BY38 KARLSÖDJ October







— Mean 2001-2015 St.Dev. • 2016-10-16





STATION SLÄGGÖ SURFACE WATER (0-10m)

Annual Cycles

— Mean 2001-2015 St.Dev. • 2016




OXYGEN IN BOTTOM WATER (depth >= 50 m)



Vertical profiles SLÄGGÖ October

— Mean 2001-2015 St.Dev. • 2016-10-14



STATION N14 FALKENBERG SURFACE WATER (0-10m)

Annual Cycles



— Mean 2001-2015

■ St.Dev.

● 2016



Temperature °C

Salinity psu





PO₄ µmol/l

DIN µmol/l



SiO₃ µmol/l



O₂ saturation %



OXYGEN IN BOTTOM WATER (depth >= 25 m)

O₂ ml/l

O₂ ml/l



Vertical profiles N14 FALKENBERG October

— Mean 2001-2015 St.Dev. • 2016-10-13

