

Oxygen Survey in the Baltic Sea 2013

- Extent of Anoxia and Hypoxia, 1960-2013



Front: Oxygen samples taken from increasing depths ready for Winkler analysis. Strong yellow colour indicate high oxygen concentrations and transparent low or none oxygen. Photo by Philip Axe.

REPORT OCEANOGRAPHY No. 49, 2013

**Oxygen Survey in the Baltic Sea 2013
- Extent of Anoxia and Hypoxia, 1960-2013**

Martin Hansson & Lars Andersson

Swedish Meteorological and Hydrological Institute, Göteborg, Sweden

Summary

A climatology atlas of the oxygen situation in the deep water of the Baltic Sea was first published in 2011 in SMHI Report Oceanography No 42. Since 2011, annual updates have been made as additional data have been reported to ICES. In this report the results for 2012 have been updated and the preliminary results for 2013 are presented. Oxygen data from 2013 have been collected during the annual Baltic International Acoustic Survey (BIAS) and from national monitoring programmes with contributions from Sweden, Poland, Finland, Estonia Lithuania, and Latvia.

For the autumn period, August to October, each profile in the dataset was examined for the occurrence of hypoxia (oxygen deficiency) and anoxia (total absence of oxygen). The depths of onset of hypoxia and anoxia were then interpolated between sampling stations producing two surfaces representing the depth at which hypoxic and anoxic conditions are found. The volume and area of hypoxia and anoxia have been calculated and the results have then been transformed to maps and diagrams to visualize the annual autumn oxygen situation during the analysed period.

The updated results for 2012 and the preliminary results for 2013 show that the extreme oxygen conditions in the Baltic Proper after the regime shift in 1999 continue. Both the areal extent and the volume of anoxia have, after the regime shift, been constantly elevated to levels only observed occasionally before the regime shift. In the Baltic Proper, Gulf of Finland and Gulf of Riga ~15% of the bottom areas was affected by anoxia and ~30% by hypoxia during 2013.

Sammanfattning

En klimatologisk atlas av syresituationen i Östersjöns djupvatten publicerades 2011 i SMHIs Report Oceanography No 42. Sedan 2011 har rapporten uppdateras årligen då kompletterande data från länder runt Östersjön har rapporterats till ICES. I denna rapport har resultaten från 2012 uppdaterats. De preliminära resultaten för 2013 baseras på data insamlade under Baltic International Acoustic Survey (BIAS) och nationell miljöövervakning med bidrag från Finland, Estland, Lettland, Litauen, Polen och Sverige.

Förekomsten av hypoxi (syrebrist) och anoxi (helt syrefria förhållanden) under höstperioden, augusti till oktober, har undersökts i varje mätprofil. Djupet där hypoxi eller anoxi först påträffas i en profil har interpolerats mellan provtagningsstationer och kombinerats med en djupdatabas för beräkning av utbredning och volym av hypoxiska och anoxiska förhållanden. Resultaten har överförts till kartor och diagram för att visualisera syresituationen i Östersjöns djupvatten.

Resultaten för 2012 och de preliminära resultaten för 2013 visar att de extrema syreförhållanden som observerats i Egentliga Östersjön efter regimskiftet 1999 fortsätter. Utbredningen av anoxi fortsätter att vara konstant förhöjd till nivåer som bara observerats i Östersjön vid enstaka år före 1999. Under 2013 beräknas ~15 % av bottenarna i Egentliga Östersjön, Finska viken och Rigabukten vara påverkade av anoxiska förhållanden och ~30% av hypoxi.

Table of contents

1	Background	1
2	Data	1
3	Method	2
4	Result	3
5	Discussion	4
6	Conclusions	5
7	Acknowledgement	6
8	References.....	6
Appendix 1 – Temperature, salinity and oxygen at BY15, Eastern Gotland Basin, 1960-2013		8
Appendix 2 - Anoxic and hypoxic areas in the Baltic Sea, 2012-2013 (The complete time series can be found in RO report 42).....		8

1 Background

The Baltic Sea is a sensitive sea area. The region is characterised by its natural formation as an enclosed estuary with high freshwater input and restricted access to oceanic high saline water. The stratification and fjord like conditions, in combination with eutrophication and other factors, form the basis for a problematic oxygen situation in the deep water.

Hypoxia is a condition that occurs when dissolved oxygen falls below the level needed to sustain most animal life. The concentration at which various animals are affected varies, but generally effects start to appear when oxygen drops below 2.8-3.4 ml/l (4- 4.8 mg/l) and acute hypoxia is usually defined between 1.4 – 2.1 ml/l (2-3 mg/l) [Rabalais, 2001; Diaz & Rosenberg, 1995; Aertebjerg et al. 2003, Swedish EPA, 2007]. It has also been shown that Baltic cod eggs need at least 2 ml/l oxygen for successful development [MacKenzie et al., 2000; Nissling, 1994; Plikshs et al., 1993; U.S. EPA, 2003; U.S. EPA, 2000,]. In this report the limit for hypoxia is set to 2.0 ml/l.

Anoxic conditions are characterised by the total absence of oxygen. When all oxygen is consumed by microbial processes hydrogen sulphide (H₂S) is formed, which is toxic for all higher marine life. Anoxic conditions lead to release of phosphate and silicate from the sediments to the water column, which, due to vertical mixing, can reach the surface layer and the photic zone. High concentrations of phosphate favour phytoplankton growth, especially cyanobacteria in the Baltic Sea during summer.

In this report time series of the bottom areal extent and water volume of anoxic and hypoxic autumn conditions of the Baltic Proper, including the Gulf of Finland and the Gulf of Riga, are presented for the period 1960 to 2013. The time series was first published in 2011 and the results have been updated annually as new additional data have become available at ICES¹. In the report from 2011 a distinct regime shift in the oxygen situation in the Baltic Proper was found in 1999. During the first regime, 1960-1999, hypoxia affected large areas while anoxic conditions were found only in minor deep areas. After the regime shift in 1999, both areal extent and volume of anoxia have been constantly elevated to levels that only occasionally have been observed before 1999. [Hansson et. al, 2011]

The report includes maps of bottom areas affected by oxygen deficiencies which can be used as a climatological atlas describing the historical development and the present oxygen situation in the Baltic Proper.

2 Data

The results for 2013 are preliminary and based on oxygen data collected during the annual Baltic International Acoustic Survey (BIAS) complemented by national and regional monitoring programmes with contributions from Estonia, Finland, Latvia, Lithuania, Poland and Sweden. These data have been subject to initial quality control only (quality assured laboratory procedures; timing and position checks; range checking). The time series and the results presented for 2013 will be updated when additional data are reported to ICES in 2014.

Data from the BIAS cruises are well suited for concurrent oxygen surveys because of the vast spatial distribution of sampling occasions and since cruises are performed by different countries, most parts of the offshore Baltic Proper are monitored.

¹ ICES Dataset on Ocean Hydrography. The International Council for the Exploration of the Sea, Copenhagen 2009.

The surveys are also performed during the autumn period (September/October) when the oxygen situation usually is most severe. Hence, this is an essential contribution of oxygen data, complementing the regular national and regional monitoring performed monthly at fixed stations.

3 Method

To process the dataset a few station profiles had to be filtered out: for example when data was missing in the deep water or when questionable data were found.

For the autumn period, August to October, each vertical profile including at least three data points, was examined for the occurrence of hypoxia (<2 ml/l) and anoxia (<0 ml/l). To find the depth of the onset of hypoxia and anoxia in each vertical profile, interpolation between discrete measurements in the profile was used. If hypoxia or anoxia was not found in the profile, the two deepest measurements in the profile were used to linearly extrapolate the oxygen concentration down towards the bottom. If two or more profiles were found at the same position an average profile was calculated for that position.

The depths of the onset of hypoxia and anoxia were gridded with linear interpolation (Delaunay triangulation) between sampling stations, producing a surface representing the depth at which hypoxic and anoxic conditions are found. The surface has then been compared with bathymetry data, [Seifert, 2001] see Figure 1, to exclude profiles where the hypoxic and anoxic depths were greater than the actual water depth. After filtering the results, the affected area and volume of hypoxia and anoxia have been calculated for each year.




Figure 1. Bathymetry [Seifert, 2001] of the south Baltic Sea and pathway of inflowing deep water during inflows. The Baltic Proper includes the Arkona Basin, the Bornholm Basin, the Gulf of Gdansk and the Eastern-, Western- and Northern Gotland Basin [Fonselius, 1995].

Areal extent and volumes are presented in relation to the area and volume of the Baltic Proper, including the Gulf of Finland and the Gulf of Riga, see Figure 1[Fonselius, 1995].

4 Result

The areal extent and volume affected by hypoxia and anoxia during the period 1960 - 2013 are presented in Figures 2 and 3 respectively. Maps presenting bottom areas affected by hypoxia and anoxia during the autumn period 2012 and 2013 can be found in Appendix 2. The mean areal extent and volume affected by hypoxia and anoxia before and after the regime shift in 1999 (see Background section or [Hansson et. al, 2011]) and the preliminary results for 2013 are presented in Table 1.

Table 1. Mean and maximum areal extent and volume of anoxia and hypoxia before and after the regime shift. Results are given as part (%) of the area and volume of the Baltic Proper, including the Gulf of Finland and the Gulf of Riga. Updated table from Hansson et. al., 2011 & 2012. Note that the results for 2013 are preliminary.

in %	1960 – 1998		1999 – 2012		2013	
	Hypoxi	Anoxi	Hypoxi	Anoxi	Hypoxi	Anoxi
Mean Areal extent	22	5	28	15	29	15
Max Areal extent (Year)	27 (1968)	14 (1969)	32 (2007)	19 (2011)	-	-
Mean Volume	13	2	18	8	19	9
Max Volume (Year)	19 (1965)	8 (1969)	21 (2011)	12 (2011)	-	-




Figure 2. Areal extent of anoxic and hypoxic conditions in the Baltic Proper, Gulf of Finland and Gulf of Riga. Results from 1961 and 1967 have been removed since sufficient data from the deep basins are missing.

Water volume affected by hypoxia and anoxia





Figure 3. Volume of anoxic and hypoxic deep water in the Baltic Proper, Gulf of Finland and Gulf of Riga. Results from 1961 and 1967 have been removed since sufficient data from the deep basins are missing.

5 Discussion

Results for 2012

When the preliminary results for 2012 were updated the areal extent and volume were only slightly adjusted. Additional anoxic areas were found in the outer parts of the Gulf of Finland and hypoxia affected large areas in the south-eastern parts of the Baltic Proper. The results for 2012 follow the oxygen development that has prevailed since the beginning of 2000s. Positive effects of the relatively large inflow in November/December 2011, $\sim 50 \text{ km}^3$ through the Sound [SMHI, 2013], could be seen in the southern Baltic Proper and in the southern parts of the eastern Gotland Basin but not in the deeper central parts.

The cumulative flow out of the Baltic Sea in 2012 was higher than average and higher than in 2011, mainly due to high precipitation. Some small inflows in February and December led to some oxygenation of the deep water in the southern Baltic. During January, northeasterly winds forced water through the Sound, out of the Baltic Proper. As a result, an inflow, through the Sound, of $\sim 35 \text{ km}^3$ during February improved the oxygen conditions in the Arkona- and the Bornholm Basins. The high precipitation and large flows in the watercourses during autumn resulted in continuous outflows through the Sound. At Christmas and the following period water levels rose along the Swedish west coast and gave rise to an inflow of $\sim 30 \text{ km}^3$ through the Sound, that again improved the oxygen situation in the southwestern part of the Baltic Proper, but was too small to affect the central parts.

Preliminary results for 2013

The preliminary results for 2013 are similar to the mean conditions for the period (1999-2012) after the regime shift. Though, in comparison with 2012, the area affected by anoxia in the western Gotland basin has increased and anoxia was again present in the Bornholm Basin. Hypoxic conditions were now also found in the Gulf of Finland.

During 2013, inflows through the Sound were registered in January/February $\sim 30 \text{ km}^3$ and together with the minor inflow in December 2012 only a temporary improvement could be seen in the Hanö Bight and in the Bornholm Basin before the oxygen concentrations again dropped below 2 ml/l to almost 0 ml/l. It was first after the inflow in October ($\sim 40 \text{ km}^3$) that a real improvement could be seen when concentrations increased from 0 ml/l to over 2 ml/l. At the time of writing, the storm Sven had just passed over the southern Baltic region. This storm resulted in high sea levels in south-eastern Kattegat and in the Sound, and in mid-December an inflow was ongoing which had reached $\sim 20 \text{ km}^3$.

Further into the southeastern Baltic Proper and the southern Eastern Gotland Basin conditions during 2013 have been stable below 2 ml/l, with the exception of two short pulses of oxygenated water, but no hydrogen sulphide have been found in this area.

In the central deep parts of the Eastern Gotland Basin and in the Western Gotland Basin the stagnation continues. At station BY15 in the Gotland deep below 200m, see Appendix 1, there are no signs, in salinity, temperature or oxygen concentrations, of any inflow after 2007. Hence, no inflow that has occurred in later years has been sufficiently large and dense to replace the deep water in the Eastern Gotland basin. Instead, the inflows that have occurred have been set at intermediate depths, between 100-175 meters. Consequently, since most likely no inflows reach the Western Gotland Basin, hypoxia and anoxia are now found at shallow depths. Hypoxia from 60 meters depth and anoxia from 67 meters depth.

The latest major inflow occurred in winter 2003-2004 and before that in 1993.

6 Conclusions

- The extreme oxygen condition in the Baltic Proper continues undiminished during 2013. The areal extent and the volume of anoxia have since the regime shift in 1999 been constantly elevated and there are no signs that inflows, weakening of the stratification or other factors have improved the oxygen conditions in the central deeper parts.
- Anoxic conditions affected $\sim 15 \%$ of the bottom areas in the Baltic Proper, including the Gulf of Finland and the Gulf of Riga and $\sim 30\%$ suffered from hypoxia during the autumn of 2013. The largest areas of anoxic conditions, $\sim 20 \%$ were found in 2011.
- The inflows that occurred at the end of 2011 and a couple of small inflows that occurred during 2012 have improved the oxygen conditions temporarily in Arkona, Hanö Bight, in the Bornholm Basin and the southern Eastern Gotland Basin. But they did not improve the conditions in the deeper central parts of the Baltic Proper.
- In 2013 an inflow in October improved the oxygen conditions remarkably in the Hanö Bight and the Bornholm Basin. The effects of another inflow in December have still not been registered in monitoring data.

7 Acknowledgement

Data for updating the 2012 results were collected at the excellent web service at the International Council for the Exploration of the Sea (ICES), making the ICES Dataset on Ocean Hydrography available.

Many thanks to:

Tycjan Wodzinowski, Department of Fishery Oceanography and Marine Ecology, Sea Fisheries Institute, Poland. Fausta Svecova & Maris Plikshs, Institute of Food Safety, Animal Health and Environment, BIOR, Latvia. Marijus Spegys, Fisheries Service under the Ministry of Agriculture, Lithuania. Aiste Kubiliute, Head of Data Management and Programmes Division, Marine Research Department, Environment Protection Agency, Lithuania. Riikka Hietala, Head of Marine Technology Services, Finnish Meteorological Institute and Tiit Raid, Estonian Marine Institute, University of Tartu, Estonia for their cooperation regarding exchange of oxygen data and many thanks to Swedish University of Aquatic Science, Department of Aquatic Resources for good cooperation regarding the SMHI oxygen survey onboard R/V Dana during the BIAS cruise.

8 References

Aertebjerg, G., Carstensen, J., Axe, P., Druon, J.-N. & Stips, A., 2003: The oxygen Depletion Event in the Kattegat, Belt Sea and Western Baltic. Baltic Sea Environment Proceedings No. 90. Helsinki Commission Baltic Marine Environment Protection Commission. ISSN 0357-2994.

Diaz, R. J. & Rosenberg, R., 1995: Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna, *Oceanogr. Mar. Bio. Ann. Rev.*, 33, 245-303.

Fonselius, S., 1995: Västerhavets och Östersjöns Oceanografi. ISBN 91-87996-07-3.

Hansson, M., Andersson, L. & Axe, P., 2011: Areal Extent and Volume of Anoxia and Hypoxia in the Baltic Sea, 1960-2011, Report Oceanography no 42, ISSN: 0283-1112.

MacKenzie, B., Hinrichsen, H.H., Plikshs, M., Wieland, K., Zezera, A.S., 2000: Quantifying environmental heterogeneity : habitat size necessary for successful development of cod *Gadus morhua* eggs in the Baltic Sea. *Marine Ecology - Progress Series*, vol: 193, pages: 143-156.

Nausch, G., Feistel, R., Umlauf, L., Mohrholz, V., Nagel, K., Siegel, H., 2012: Hydrographisch-chemische Zustandseinschätzung der Ostsee 2011, *Meereswissenschaftliche Berichte MARINE SCIENCE REPORTS* No. 86. Leibniz- Institut für Ostseeforschung Warnemünde.

Nissling, A., 1994: Survival of eggs and yolk sac larvae of Baltic cod (*Gadus morhua*) at low oxygen levels in different salinities. *ICES Marine Science Symposium* 198:626-631.

Plikshs, M., Kalejs, M. & Grauman, G., 1993: The influence of environmental conditions and spawning stock size on the year-class strength of the Eastern Baltic cod. *ICES CM* 1993/J:22.

Rabalais, N. N. & Eugene, R., Turner (Editors), 2001: Coastal and Estuarine Studies, Coastal Hypoxia, Consequences for living resources and ecosystems. American Geophysical Union. ISBN 0-87590-272-3.

SMHI, 2013: Cruise report archive: <http://www.smhi.se/en/theme/marine-environment-2-885>.
Updated: December, 2013.

SMHI, 2013: Accumulated inflow through the Öresund. URL:
http://www.smhi.se/hfa_coord/BOOS/Oresund.html

Swedish EPA, 2007: Bedömningsgrunder för kustvatten och vatten i övergångszonen, Bilaga B till handboken 2007:4, Naturvårdsverket, ISBN 978-91-620-0149-0.


T. Seifert, F. Tauber, B. Kayser: 2001: A high resolution spherical grid topography of the Baltic Sea – 2nd edition, Baltic Sea Science Congress, Stockholm 25-29. November 2001, Poster #147.

U.S. EPA, 2003: Ambient Water Quality Criteria for Dissolved Oxygen, Water Clarity and Chlorophyll a for the Chesapeake Bay and Its Tidal Tributaries, U.S. Environmental Protection Agency.

U.S. EPA, 2000: Ambient Aquatic Life Water Quality Criteria for Dissolved Oxygen (Saltwater): Cape Cod to Cape Hatteras, U.S. Environmental Protection Agency, EPA-822-R-00-012.

Appendix 1 – Temperature, salinity and oxygen at BY15, Eastern Gotland Basin, 1960-2013


BY15 (GOTLAND DEEP) 240m



Appendix 2 - Anoxic and hypoxic areas in the Baltic Sea, 2012-2013


(The complete time series can be found in RO report 42)

Extent of hypoxic & anoxic bottom water, Autumn 2012



Created:
December 2013

Extent of hypoxic & anoxic bottom water, Autumn 2013



Created:
December 2013

9 SMHI Publications

SMHI publish seven report series. Three of these, the R-series, are intended for international readers and are in most cases written in English. For the others the Swedish language is used.

Name of the series	Published since
RMK (Report Meteorology and Climatology)	1974
RH (Report Hydrology)	1990
RO (Report Oceanography)	1986
METEOROLOGI	1985
HYDROLOGI	1985
OCEANOGRAFI	1985
KLIMATOLOGI	2009

Earlier issues published in RO

- | | |
|--|--|
| <p>1 Lars Gidhagen, Lennart Funkquist and Ray Murthy (1986)
Calculations of horizontal exchange coefficients using Eulerian time series current meter data from the Baltic Sea.</p> <p>2 Thomas Thompson (1986)
Ymer-80, satellites, arctic sea ice and weather.</p> <p>3 Stig Carlberg et al (1986)
Program för miljö kvalitetsövervakning - PMK.</p> <p>4 Jan-Erik Lundqvist och Anders Omstedt (1987)
Isförhållandena i Sveriges södra och västra farvatten.</p> <p>5 Stig Carlberg, Sven Engström, Stig Fonselius, Håkan Palmén, Eva-Gun Thelén, Lotta Fyrberg och Bengt Yhlen (1987)
Program för miljö kvalitetsövervakning - PMK. Utsjöprogram under 1986.</p> | <p>6 Jorge C. Valderama (1987)
Results of a five year survey of the distribution of UREA in the Baltic sea.</p> <p>7 Stig Carlberg, Sven Engström, Stig Fonselius, Håkan Palmén, Eva-Gun Thelén, Lotta Fyrberg, Bengt Yhlen och Danuta Zagradkin (1988).
Program för miljö kvalitetsövervakning - PMK. Utsjöprogram under 1987</p> <p>8 Bertil Håkansson (1988)
Ice reconnaissance and forecasts in Storfjorden, Svalbard.</p> <p>9 Stig Carlberg, Sven Engström, Stig Fonselius, Håkan Palmén, Eva-Gun Thelén, Lotta Fyrberg, Bengt Yhlen, Danuta Zagradkin, Bo Juhlin och Jan Szaron (1989)
Program för miljö kvalitetsövervakning - PMK. Utsjöprogram under 1988.</p> <p>10 L. Fransson, B. Håkansson, A. Omstedt och L. Stehn (1989)
Sea ice properties studied from the</p> |
|--|--|

- ice-breaker Tor during BEPERS-88.
- 11 Stig Carlberg, Sven Engström, Stig Fonselius, Håkan Palmén, Lotta Fyrberg, Bengt Yhlen, Bo Juhlin och Jan Szaron (1990)
Program för miljö kvalitetsövervakning - PMK. Utsjöprogram under 1989.
- 12 Anders Omstedt (1990)
Real-time modelling and forecasting of temperatures in the Baltic Sea.
- 13 Lars Andersson, Stig Carlberg, Elisabet Fogelqvist, Stig Fonselius, Håkan Palmén, Eva-Gun Thelén, Lotta Fyrberg, Bengt Yhlen och Danuta Zagradkin (1991)
Program för miljö kvalitetsövervakning – PMK. Utsjöprogram under 1989.
- 14 Lars Andersson, Stig Carlberg, Lars Edler, Elisabet Fogelqvist, Stig Fonselius, Lotta Fyrberg, Marie Larsson, Håkan Palmén, Björn Sjöberg, Danuta Zagradkin, och Bengt Yhlen (1992)
Haven runt Sverige 1991. Rapport från SMHI, Oceanografiska Laboratoriet, inklusive PMK - utsjöprogrammet. (The conditions of the seas around Sweden. Report from the activities in 1991, including PMK - The National Swedish Programme for Monitoring of Environmental Quality Open Sea Programme.)
- 15 Ray Murthy, Bertil Håkansson and Pekka Alenius (ed.) (1993)
The Gulf of Bothnia Year-1991 - Physical transport experiments.
- 16 Lars Andersson, Lars Edler and Björn Sjöberg (1993)
The conditions of the seas around Sweden. Report from activities in 1992.
- 17 Anders Omstedt, Leif Nyberg and Matti Leppäranta (1994)
A coupled ice-ocean model supporting winter navigation in the Baltic Sea.
Part 1. Ice dynamics and water levels.
- 18 Lennart Funkquist (1993)
An operational Baltic Sea circulation model. Part 1. Barotropic version.
- 19 Eleonor Marmefelt (1994)
Currents in the Gulf of Bothnia. During the Field Year of 1991.
- 20 Lars Andersson, Björn Sjöberg and Mikael Krysell (1994)
The conditions of the seas around Sweden. Report from the activities in 1993.
- 21 Anders Omstedt and Leif Nyberg (1995)
A coupled ice-ocean model supporting winter navigation in the Baltic Sea.
Part 2. Thermodynamics and meteorological coupling.
- 22 Lennart Funkquist and Eckhard Kleine (1995)
Application of the BSH model to Kattegat and Skagerrak.
- 23 Tarmo Köuts and Bertil Håkansson (1995)
Observations of water exchange, currents, sea levels and nutrients in the Gulf of Riga.
- 24 Urban Svensson (1998)
PROBE An Instruction Manual.
- 25 Maria Lundin (1999)
Time Series Analysis of SAR Sea Ice Backscatter Variability

and its Dependence on Weather Conditions.

- 26 Markus Meier¹, Ralf Döscher¹, Andrew, C. Coward², Jonas Nycander³ and Kristofer Döös³ (1999). RCO – Rossby Centre regional Ocean climate model: model description (version 1.0) and first results from the hindcast period 1992/93.

¹ Rossby Centre, SMHI ² James Rennell Division, Southampton Oceanography Centre, ³ Department of Meteorology, Stockholm University

- 27 H. E. Markus Meier (1999) First results of multi-year simulations using a 3D Baltic Sea model.
- 28 H. E. Markus Meier (2000) The use of the $k - \epsilon$ turbulence model within the Rossby Centre regional ocean climate model: parameterization development and results.
- 29 Eleonor Marmefelt, Bertil Håkansson, Anders Christian Erichsen and Ian Sehested Hansen (2000) Development of an Ecological Model System for the Kattegat and the Southern Baltic. Final Report to the Nordic Councils of Ministers.
- 30 H.E Markus Meier and Frank Kauker (2002). Simulating Baltic Sea climate for the period 1902-1998 with the Rossby Centre coupled ice-ocean model.
- 31 Bertil Håkansson (2003) Swedish National Report on Eutrophication Status in the Kattegat and the Skagerrak OSPAR ASSESSMENT 2002

- 32 Bengt Karlson & Lars Andersson (2003) The Chattonella-bloom in year 2001 and effects of high freshwater input from river Göta Älv to the Kattegat-Skagerrak area
- 33 Philip Axe and Helma Lindow (2005) Hydrographic Conditions Around Offshore Banks
- 34 Pia M Andersson, Lars S Andersson (2006) Long term trends in the seas surrounding Sweden. Part one - Nutrients
- 35 Bengt Karlson, Ann-Sofi Rehnstam-Holm & Lars-Ove Loo (2007) Temporal and spatial distribution of diarrhetic shellfish toxins in blue mussels, *Mytilus edulis* (L.), at the Swedish West Coast, NE Atlantic, years 1988-2005
- 36 Bertil Håkansson Co-authors: Odd Lindahl, Rutger Rosenberg, Pilip Axe, Kari Eilola, Bengt Karlson (2007) Swedish National Report on Eutrophication Status in the Kattegat and the Skagerrak OSPAR ASSESSMENT 2007
- 37 Lennart Funkquist and Eckhard Kleine (2007) An introduction to HIROMB, an operational baroclinic model for the Baltic Sea
- 38 Philip Axe (2008) Temporal and spatial monitoring of eutrophication variables in CEMP

- 39 Bengt Karlson, Philip Axe, Lennart Funkquist, Seppo Kaitala, Kai Sørensen (2009)
Infrastructure for marine monitoring and operational oceanography
- 40 Marie Johansen, Pia Andersson (2010)
Long term trends in the seas surrounding Sweden
Part two – Pelagic biology
- 41 Philip Axe, (2012)
Oceanographic Applications of Coastal Radar
- 42 Martin Hansson, Lars Andersson, Philip Axe (2011)
Areal Extent and Volume of Anoxia and Hypoxia in the Baltic Sea, 1960-2011
- 43 Philip Axe, Karin Wesslander, Johan Kronsell (2012)
Confidence rating for OSPAR COMP
- 44 Germo Väli, H.E. Markus Meier, Jüri Elken (2012)
Simulated variations of the Baltic Sea halocline during 1961-2007
- 45 Lars Axell (2013)
BSRA-15: A Baltic Sea Reanalysis 1990-2004
- 46 Martin Hansson, Lars Andersson, Philip Axe, Jan Szaron (2013)
Oxygen Survey in the Baltic Sea 2012 - Extent of Anoxia and Hypoxia, 1960 -2012
- 47 C. Dieterich, S. Schimanke, S. Wang, G. Väli, Y. Liu, R. Hordoir, L. Axell, A. Höglund, H.E.M. Meier (2013)
Evaluation of the SMHI coupled atmosphere-ice-ocean model RCA4-NEMO
- 48 R. Hordoir, B. W. An, J. Haapala, C. Dieterich, S. Schimanke, A. Höglund and H.E.M. Meier (2013)
BaltiX V 1.1 : A 3D Ocean Modelling Configuration for Baltic & North Sea Exchange Analysis



Swedish Meteorological and Hydrological Institute
SE 601 76 NORRKÖPING
Phone +46 11-495 80 00 Telefax +46 11-495 80 01

ISSN 0283-1112