

Simulated distribution of colored dissolved organic matter in the Baltic Sea

Ulrike Löptien, H.E. Markus Meier



Front.
Yellowisch river water with light reflection
Foto: U. Löptien

OCEANOGRAPHI Nr 109, 2011

**Simulated distribution of colored dissolved
organic matter in the Baltic Sea**

Ulrike Löptien, H.E. Markus Meier

Summary

The Baltic Sea is optically a multi-componental water and has exceedingly high levels of colored dissolved organic matter (CDOM, also referred as yellow matter, gilvin or gelbstoff). CDOM is a complex mixture of chemical bonds originating from the decay of photosynthetically produced organic matter. It influences the aquatic light field substantially. A quantitative description of the dynamics and variability is often required to predict accurately light penetration and hereafter e.g. primary production.

The present study is a first attempt to include CDOM into a Baltic Sea model. The model integrations are based on a fixed concentration in the 30 major rivers. In the absence of comprehensive measurements, a river inflow proportional to total organic carbon is assumed. Since origin and fate are still a matter of discussion, we test various decay rates of CDOM and compare the results with satellite observations. Best results are obtained when assuming a light dependent decay, compared to a temperature or time dependent decay. Treating CDOM as a conservative tracer does not lead to satisfactory results.

Sammanfattning

Vattnet i Östersjön är optiskt sett uppdelat i många beståndsdelar och har ytterst höga halter av färgade lösta organiska ämnen (colored dissolved organic matter, CDOM), även kallade för gulämnen, gilvin eller gelbstoff. CDOM är en komplex blandning av kemiska bindningar skapade vid nedbrytningen av fotosyntetiskt producerat organiskt material och det har en avsevärd påverkan på ljusfältet i vattnet. En kvantitativ beskrivning av dynamiken och variabiliteten behövs ofta för att få en riktig beskrivning av ljusets nedträngning och där efter av t.ex. primärproduktionen.

Denna studie är ett första försök att inkludera CDOM i en modell för Östersjön. Modellförsöken baseras på en fix koncentration i de 30 största floderna. I brist på omfattande mätningar antas en flödstillförsel som är proportionell mot totalt organiskt kol. Eftersom ursprunget och ödet fortfarande är föremål för diskussioner testas vi olika hastigheter av CDOM sönderfall och jämför resultaten med satellitobservationer. Jämfört med temperatur eller tidsberoende sönderfall uppnåddes bäst resultat med ett ljusberoende sönderfall. Att behandla CDOM som ett konservativt spårämne leder inte till tillfredsställande resultat.

1 Introduction

The Baltic Sea is located in central Europe and about 85 million people are living in its drainage basin (e.g. [Leppäranta and Myrberg \(2009\)](#)). This optically multi-componental water comprises extremely high levels of colored dissolved organic matter (CDOM, also referred as yellow matter, gilvin or gelbstoff), contained by the low exchange of Baltic waters with the North Sea ([Jerlov \(1968\)](#), [Siegel et al. \(2005\)](#), [Kratzer and Tett \(2009\)](#)). CDOM is, by definition, the part of dissolved organic matter which is capable to absorb light. It is a complex mixture of chemical bonds originating from the decay of photosynthetically produced organic matter and can play a substantial role in the biogeochemistry of natural waters merely through its influence on the aquatic light field (e.g. [Bidigare and Ondrusek \(1997\)](#), [Blough et al. \(1993\)](#), [Blough \(1996\)](#), [Moran and Zepp \(2004\)](#), [Whithead et al. \(1997\)](#)). Thus, a quantitative description of the dynamics and variability of CDOM is often required to predict accurately light penetration and hereafter e.g. primary production.

The origin and fate of CDOM is still a matter of discussion ([del Vecchio and Blough \(2006\)](#)). [Siegel et al. \(2002\)](#) analysed global satellite data of colored detrital and dissolved materials (CDM). Here, detrital particulates are assumed to make only a small contribution to CDM. The authors point out the different origin of coastal and open ocean dissolved organic matter. Near coastal CDOM, like in the Baltic Sea, is mainly regulated by land-ocean interactions and originates mainly from land drainage and river runoff. Entering the ocean, CDOM is mixed vertically, advected and decomposed (e.g. by photobleaching, flocculation), while new production is assumed to play a minor role. The importance of a CDOM sink compared to other effects (e.g. dilution) is still under discussion. In the absence of a sink of terrestrial CDOM, it would take about 35,000 years for the ocean to reach average river CDOM absorption when assuming the present-day total annual river flow. The existing large gradient between fresh water and 'blue water' implies the presence of a loss mechanism acting on far shorter time scales ([Blough and Del Vecchio \(2002\)](#)). It has been shown by a large number of field studies that photochemistry alone can act as a substantial sink of terrestrial CDOM ([Mopper and Kieber \(2002\)](#)). Also, laboratory experiments on (optically) thin natural waters have shown absorption losses

greater than 50% over irradiation periods of several hundred to thousand hours using sources approximating the surface solar spectrum. The time scale of these losses, on the order of weeks or month under natural irradiation conditions, is consistent with field studies by [Vodacek et al. \(1997\)](#) and [Nelson et al. \(1998\)](#) in the North Atlantic and the Saragossa Sea. On the other hand, in the presence of high CDOM-concentrations photobleaching is restricted to a very thin surface layer due to the high CDOM-absorption ([Blough and Del Vecchio \(2002\)](#)) and the net effect might be negligible. Some studies of the Baltic Sea - North Sea transition zone even treat CDOM as conservative tracer to identify the origin of water masses ([Højerslev et al. \(1996\)](#), [Sedmon et al. \(2010\)](#)). This assumption is based on the close relation between salinity and CDOM reported e.g. by [Ferrari et al. \(1998\)](#), [Kowalczyk et al. \(1998\)](#), [Kowalczyk et al. \(2006\)](#).

The present study is a first attempt to include CDOM in a Baltic Sea model. At this, we assume various decay rates of CDOM and compare the results with satellite observations. Also, we are testing the hypothesis that CDOM can be regarded as conservative tracer which is of great interest for mixing studies of water masses.

2 Methods and model

2.1 The ocean model

The oceanic component consists of the three-dimensional 'Rossby Centre Regional Ocean model' (RCO). The model setup is described in detail e.g. by [Meier et al. \(2003\)](#) and was successfully used in various climate related studies ([Meier and Kauker \(2003\)](#), [Meier et al. \(2004\)](#), [Meier \(2005\)](#)). RCO was originally a regionalized version of the OCCAM-model ([Webb et al. \(1997\)](#)). The model domain covers the Baltic Sea including Kattegat. The horizontal resolution used here is 2nm using 41 vertical layers. As surface boundary conditions air temperature, wind, cloud cover, sea level pressure, humidity and precipitation are required while surface restoring is not applied. Since as well common reanalysis data sets as well as global climate models have their limitations in regional detail, we use regional climate models with

a limited model domain and higher resolution to downscale coarser resolved data sets. Here, we use the Rossby Centre Regional Atmosphere model (RCA) at 25 km horizontal resolution (Jones et al. (2004), [Samuelsson et al. \(2011\)](#)). The boundary conditions for the atmospheric model are taken from ERA40 reanalysis data (Uppala et al. (2005)) .

2.2 Colored dissolved organic matter

CDOM is induced into the ocean model by assuming a certain concentration in the 30 major rivers. River runoff data are taken from [Bergström and Carlsson \(1994\)](#) and updated with results from a large-scale hydrological model ([Graham \(2004\)](#)). The crossover from observations to the hydrological model causes for some rivers a change in variability while the mean values remain basically unchanged. In the absence of comprehensive measurements of yellow substances (or dissolved organic carbon) we assume a river inflow proportional to total organic carbon (TOC). Measurements were provided by [Humborg et al. \(2008\)](#). For five of the rivers used by our model no measurements are provided and we assume basin mean concentrations. The position of the considered rivers is depicted in Fig. 1. The size of the dots is proportional to the inflow of yellow substances per river and results from the product of volume flow times concentration of TOC. The differences between the rivers are dominated by the volume of water inflow per river while, in general, the mean TOC concentration per river differs by less than a factor of two. Our modeling approach assumes, that new production of CDOM plays a minor role and that the North Sea has a negligible content of yellow substances. The latter assumption is in close agreement with the results presented by [Højerslev et al. \(1996\)](#). Under this assumptions we performed a set of experiments based on different decay rates of CDOM. All experiments are listed in Table 1.

In the first five experiments, we assume different decay rates from no decay and light dependent decay with a maximum half live time 240, 480, 960 and 1920 hours. The decay rates are calculated for every grid box depending on the downward radiative flux in RCO which is parametrized by two exponential functions following [Paulson and Simpson \(1977\)](#). Maximal decay is reached in the surface grid box at an in-




Figure 1: *Position of the rivers used in the model setup. The size of the dots is proportional to the amount of yellow substances discharged by the particular river. The amount of yellow substances results from water supply times concentration.*

Exp.	Description	Min. half life time
NODECAY	no decay of yellow substances	-
LIGHT240	light dependent decay	240 hours
LIGHT480	light dependent decay	480 hours
LIGHT960	light dependent decay	960 hours
LIGHT1920	light dependent decay	1920 hours
ZERF1920	constant decay	1920 hours
TEMP960	temperature dependent decay	1920 hours

Table 1: *Acronyms for model experiments discussed in the text (first column), a short description and the considered half life time.*

coming radiation of 900 W/m^2 . Below this maximum this decay decreases linearly with the available light. The amplitude of the decay is given by the fraction of the effective mean radiation and the maximal mean radiation in the upper grid box. Two additional comparative experiments assume a constant and a temperature dependent decay of CDOM. The temperature dependent decay is designed such that the decay is maximal at 20°C and above. At lower temperatures (TEMP) the decay decreases by a factor of $TEMP/20$. A more systematic approach to test different decay rates is not possible due to the high complexity and long run-times of the ocean model. Following Meier (2005) we performed a 90 year spinup by repeating present day forcing (1962-2007) twice. Additionally, discharge experiments of the three major rivers were carried out.

2.3 Observational data

In the absence of comprehensive CDOM-measurements, we focus mainly on satellite data during the period 2003-2006 which were provided by the European Space Agency (<http://www.esa.int/esaCP/index.html>). Note, however, that satellite measurements in the Baltic Sea suffer from difficulties due to the atmospheric composition and the multi-componental water properties (Darecki (2003), Kratzer et al. (2003)).




Figure 2: *Normalized seasonal mean concentration of yellow substances as observed by satellite (MERIS).*

According to Kratzer and Tett (2009) the MERIS standard processor underestimates




Figure 3: *Normalized seasonal mean concentration of yellow substances as modeled in summer in (a) experiment NODECAY and (b) in experiment LIGHT480.*

yellow substances in the open Baltic by about 81%. This error is basically related to an off-set which implies that we have to focus on the pattern and the mean seasonal cycle rather than on absolute values. Before normalisation by dividing through the maximum value we added the offset (= 80% of the mean summer concentration of yellow substances at Gotland Deep) to obtain reasonable gradients in space and time. Note, that this is a rather rough approach which is due to the lack of more reliable data. In Fig. 2 the normalized pattern are depicted.

2.4 Model results

Comparing the spatial pattern of the modeled CDOM concentrations with satellite observations, reveals that the pattern roughly agree when considering experiment NODECAY (Fig. 3a). However, one apparent discrepancy is a somewhat too high concentration in the Gulf of Finland, too high concentrations close to the river mouth

and, apart from the regions very close to the coast, too weak spatial gradients. A light dependent decay basically enhances the spatial gradient while the overall pattern remains in principle unchanged (e.g. LIGHT480 in Fig. 3b).

Considering the large scale time behaviour shows as well an underestimation of the variability when no decay of yellow substances is assumed. Fig 4a depicts the mean seasonal cycle of CDOM in the centre of the Baltic proper (14° - 19° E, 54° - 58° N) in the experiment NODECAY in comparison to observations. In contrast to the observations, which show a pronounced minimum during the summer month, CDOM is accumulating throughout the year until mixed layer depth increases during winter and run-off is reduced at the same time (Fig. 4). The observed seasonal cycle can not be reproduced by either a constant (ZERF1920) or temperature related decay (TEMP960). While a constant decay does not change the shape of the seasonal cycle at all, but increases the amplitude, a temperature dependent decay leads to a decrease of CDOM in late summer/autumn (Fig. 4a) when water temperatures are highest. As shown in Fig. 5, similar results are obtained in the northern Baltic Sea (16° - 24° E, 60° - 70° N). The assumption of a light dependent decay gives most satisfactory results (Fig. 4b and Fig. 5b) while the results concerning the strength are somewhat contradictory. In experiment LIGHT240 the amplitude of the seasonal cycle compares best to observations (though the seasonal cycle is still somewhat too weak and the increase in the yellow substance content is a bit too slow in autumn), but the spatial gradient is too strong and a weaker light dependent decay seems likely.

3 Conclusions

In the present study, CDOM was included into a Baltic Sea model assuming a river supply proportional to TOC. At first order, we assume that new production is of minor importance and that the North Sea has a negligible content of yellow substances. Given this assumptions, we obtain a reasonable agreement with observed pattern, while some large differences occur in detail. Our attempt to model yellow substances in the Baltic Sea was most successful when assuming a linear light dependent decay

of yellow substances with a half life time of 240-480 hours under maximum light exposure. Thus, according to our finding yellow substances can not be regarded as a conservative tracer. This result is inline to the results of Opsahl and Brenner (1998) who report a loss of 75% of total dissolved ligning during 28d of light exposure at the Mississippi river plume.




Figure 4: *Seasonal cycle of yellow substances (normalized by division through the maximum value) in the Baltic Proper as modeled. (a) NODECAY (black line), ZERF1920 (green line) and TEMP960 (red line). The observed, normalized values in the same region as observed by satellite (MERIS) are depicted as red dots. (b) LIGHT240 (blue line), LIGHT480 (red line), LIGHT960 (green line) and LIGHT1920 (black line).*




Figure 5: Seasonal cycle of yellow substances (normalized by division through the maximum value) in the Bothnian Sea (16°-24° E, 60°-70° N) as modeled. (a) NODE-CAY (black line), ZERF1920 (green line) and TEMP960 (red line). The observed, normalized values in the same region as observed by satellite (MERIS) are depicted as red dots. (b) LIGHT240 (blue line), LIGHT480 (red line), LIGHT960 (green line) and LIGHT1920 (black line).

Acknowledgments

This study was performed within the projects 'Sensor Networks to Monitor Marine Environment with Particular Focus on Climate Changes' funded by the Swedish Governmental Agency for Innovation Systems (Vinnova, project-no. P29461-1) and 'Advanced modeling tool for scenarios of the Baltic Sea ECOSystem to SUPPORT decision making ECOSUPPORT' jointly funded by the Swedish Environmental Protection Agency (Naturvårdsverket, grant no. 08/381) and the European Commission within the BONUS+ program (www.bonusportal.org). The RCO model simulations were partly performed on the climate computing resources 'Ekman' and 'Vagn' that are operated by the National Supercomputer Centre (NSC) at Linköping University and the Centre for High Performance Computing (PDC) at the Royal Institute of Technology in Stockholm, respectively. These computing resources are funded by a grant from the Knut and Alice Wallenberg foundation. We are thankful for these supports.

References

- Bergström, S. and B. Carlsson, 1994: River runoff to the Baltic Sea: 1950-1990. *Ambio*, **23**, 280–287.
- Bidigare, R. R. and M. E. Ondrusek, 1997: Influence of the Orinoco River outflow on distribution of algal pigments in the Caribbean Sea. *J. Geophys. Res.*, **98**, 2259–2269.
- Blough, N. V., 1996: Photochemistry in the sea-surface microlayer. *In: P. Liss and R. Duce [eds.] The sea surface and global change. Cambridge University Press*, pp. 383–424.
- Blough, N. V. and R. Del Vecchio, 2002: Chromophoric DOM in the coastal environment. *in: Hansell and Carlson, C.A. (Ed.), Biogeochemistry of marine dissolved organic matter*, pp. 509–546.

- Blough, N. V., o. zafiriou, and j. bonilla, 1993: optical absorbtion spectra of water from the orinoco river outflow: terrestrial input of colored organic matter to the caribbean. *J. Geophys. Res.*, **98**, 2271–2278.
- Darecki, D., M.and Stramski, 2003: An evaluation of MODIS and SeaWIFS bio-optical algorithms in the Baltic Sea. *Remote Sensing of Enviroment*, **89**, 326–350.
- del Vecchio, R. and N. V. Blough, 2006: Influence of ultraviolet radiation on the chromophoric dissolved organic matter in natural waters. *Environmental UV Radiation: Impact on Ecosystems and Human Health and Predictive Models*, Springer Netherlands. DOI:10.1007/140203697316.
- Ferrari, G. M., M. D. Dowell, S. Grossi, and C. Targa, 1998: Relationship between the optical properties of chromophoric dissolved organic matter and total concentration of dissolved organic carbon in the southern baltic sea region. *Marine Chemistry*, **55**, 299–316.
- Graham, L. P., 2004: Climate change effects on river flow to the Baltic Sea. *Ambio*, **33**, 235–241.
- Højerslev, N. K., N. Holt, and T. Aarup, 1996: Optical measurements in the North Sea-Baltic sea transition zone. I On the origin of the deep water of kattegat. *Estuarine, Coastal and Shelf Science*, **16**(10), 1329–1342.
- Humborg, C., E. Smedberg, M. R. Medina, and C. M. Mörtz, 2008: Changes in dissolved silicate loads to the Baltic Sea - the effects of lakes and reservoirs. *Journal of Marine Systems*, **73**, 223–235.
- Jerlov, N., 1968: Optical oceanography. *ISBN 978-0-444-40320-9*, Elsevier.
- Jones, C., U. Willén, A. Ullerstig, and U. Hansson, 2004: The Rossby Centre Regional Atmospheric Climate Model Part I: Model climatology and performance for the present climate over europe. *Ambio*, **33**, 199–210.
- Kowalczyk, P., C. A. Sedmon, and S. Markager, 2006: Modeling apsorption by cdom in the baltic sea from season, salinity and chlorophyll. *Marine Chemistry*, **101**, 1–11.

- Kowalczyk, P., J. Ston-Egiert, R. F. Cooper, and C. Targa, 1998: Characterization of chromophoric dissolved organic matter (cdom) in the baltic sea by excitation emission matrix fluorescence spectroscopy. *Marine Chemistry*, **96**, 273–292.
- Kratzer, S., B. Håkansson, and C. Sahlin, 2003: Assessing secchi and photic zone depth in the Baltic Sea from satellite data. *Ambio*, **32**(8), 577–585.
- Kratzer, S. and P. Tett, 2009: Using bio-optics to investigate the extent of coastal waters: A Swedish case study. *Hydrobiologica*, **629**, 169–186.
- Leppäranta, M. and K. Myrberg, 2009: Physical oceanography of the Baltic Sea. *ISBN 978-3-540-79702-9, Springer*, pp. 85,296–303.
- Meier, H. E. M., 2005: Modeling the age of Baltic Sea water masses: quantification and steady state sensitivity experiments. *J Geophys Res*, **110**. Doi: 10.1029/2004JC002 607.
- Mopper, K. and D. J. Kieber, 2002: Photochemistry and the cycling of carbon, sulfur, nitrogen and phosphorus. *in: Hansell and Carlson, C.A. (Ed.), Biogeochemistry of marine dissolved organic matter*, pp. 455–507.
- Moran, M. A. and R. G. Zepp, 2004: Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter. *Limnol. Oceanogr.*, **42**, 1307–1316.
- Nelson, N. B., D. A. Siegel, and A. F. Michaels, 1998: Seasonal dynamics of colored dissolved organic matter in the Sargasso Sea. *Deep-Sea Res. I*, **45**, 931–957.
- Opsahl, S. and R. Brenner, 1998: Photochemical reactivity of dissolved lignin in river and ocean waters. *Limnol. Oceanogr.*, **43**(6), 1297–1304.
- Paulson, C. A. and J. J. Simpson, 1977: Irradiance measurements in the upper ocean. *J. Phys. Oceanogr*, **7**, 952–956.
- Samuelsson, P., C. Jones, U. Willén, S. Gollvik, and coauthors, 2011: The Rossby Centre Regional Climate Model RCA3: Model description and performance. *Tellus*, **63A**.

- Sedmon, C. A., C. L. Osburn, and T. Kragh, 2010: Tracing water mass mixing in the baltic-north sea transition zone using the optical properties of coloured dissolved organic matter. *Estuarine, Coastal and Shelf Science*, **87**, 156–162.
- Siegel, D. A., S. Maritorena, and N. B. Nelson, 2002: Global distribution and dynamics of colored dissolved and detrital organic materials. *J. Geophys. Res.*, **127**. DOI:10.1029/2001JC000965.
- Siegel, H., M. Gerth, T. Ohde, and T. Heene, 2005: Ocean colour remote sensing relevant water constituents and optical properties of the Baltic Sea. *International Journal of Remote Sensing*, **26**, 315–330.
- Uppala, S., P. W. Kallberg, A. J. Simmons, U. Andrae, V. Da Costa Bechtold, M. Fiorino, J. K. Gibson, J. Haseler, A. Hernandez, G. A. Kelly, X. Li, K. Onogi, S. Saarinen, N. Sokka, R. P. Allan, E. Andersson, K. Arpe, M. A. Balmaseda, A. C. M. Beljaars, L. van de Berg, J. Bidlot, N. Bormann, S. Caires, F. Chevallier, A. Dethof, M. Dragosavac, M. Fisher, M. Fuentes, S. Hagemann, E. Holm, B. J. Hoskins, L. Isaksen, P. A. E. M. Janssen, R. Jenne, A. P. McNally, J.-F. Mahfouf, J.-J. Morcrette, N. A. Rayner, R. W. Saunders, P. Simon, A. Sterl, K. E. Trenberth, A. Untch, D. Vasiljevic, P. Viterbo, and J. Woollen, 2005: The ERA-40 reanalysis. *Quart. J. Royal Met. Soc.*, **131**, 2961–3012.
- Vodacek, A. N., M. D. Blough, E. T. DeGrandpre, E. T. Peltzer, and R. K. Nelson, 1997: Seasonal variation of cdom and doc in the middle atlantic bight: Terrestrial inputs and photooxidation. *Limnol. Oceanogr.*, **42**, 674–686.
- Whithead, R. F., S. de Mora, S. Demers, M. Gosslin, P. Monfort, and B. Mostajir, 1997: Interactions of ultraviolet-B radiation, mixing, and biological activity on photobleaching of natural chromophoric dissolved organic matter: A mesocosm study. *Limnol. Oceanogr.*, **45**(2), 278–291.

I serien OCEANOGRAFI har tidigare utgivits:

- 1 Lennart Funkquist (1985)
En hydrodynamisk modell för spridnings-
och cirkulationsberäkningar i Östersjön
Slutrapport.
- 2 Barry Broman och Carsten Pettersson.
(1985)
Spridningsundersökningar i yttre fjärden
Piteå.
- 3 Cecilia Ambjörn (1986).
Utbyggnad vid Malmö hamn; effekter för
Lommabuktens vattenutbyte.
- 4 Jan Andersson och Robert Hillgren (1986).
SMHIs undersökningar i Öregrundsgrepen
perioden 84/85.
- 5 Bo Juhlin (1986)
Oceanografiska observationer utmed
svenska kusten med kustbevakningens
fartyg 1985.
- 6 Barry Broman (1986)
Uppföljning av sjövärmepump i Lilla
Värtan.
- 7 Bo Juhlin (1986)
15 års mätningar längs svenska kusten med
kustbevakningen (1970 - 1985).
- 8 Jonny Svensson (1986)
Vågdata från svenska kustvatten 1985.
- 9 Barry Broman (1986)
Oceanografiska stationsnät - Svenskt
Vattenarkiv.
- 10 -
- 11 Cecilia Ambjörn (1987)
Spridning av kylvatten från Öresundsverket
- 12 Bo Juhlin (1987)
Oceanografiska observationer utmed
svenska kusten med kustbevakningens
fartyg 1986.
- 13 Jan Andersson och Robert Hillgren (1987)
SMHIs undersökningar i Öregrundsgrepen
1986.
- 14 Jan-Erik Lundqvist (1987)
Impact of ice on Swedish offshore
lighthouses. Ice drift conditions in the area
at Sydostbrotten - ice season 1986/87.
- 15 SMHI/SNV (1987)
Fasta förbindelser över Öresund - utredning
av effekter på vattenmiljön i Östersjön.
- 16 Cecilia Ambjörn och Kjell Wickström
(1987)
Undersökning av vattenmiljön vid
utfyllnaden av Kockums varvsbassäng.
Slutrapport för perioden
18 juni - 21 augusti 1987.
- 17 Erland Bergstrand (1987)
Östergötlands skärgård - Vattenmiljön.
- 18 Stig H. Fonselius (1987)
Kattegatt - havet i väster.
- 19 Erland Bergstrand (1987)
Recipientkontroll vid Breviksnäs fiskodling
1986.
- 20 Kjell Wickström (1987)
Bedömning av kylvattenrecipienten för ett
kolkraftverk vid Oskarshamnsverket.
- 21 Cecilia Ambjörn (1987)
Förstudie av ett nordiskt modellsystem för
kemikaliespridning i vatten.
- 22 Kjell Wickström (1988)
Vågdata från svenska kustvatten 1986.
- 23 Jonny Svensson, SMHI/National Swedish
Environmental Protection Board (SNV)
(1988)
A permanent traffic link across the
Öresund channel - A study of the hydro-
environmental effects in the Baltic Sea.
- 24 Jan Andersson och Robert Hillgren (1988)
SMHIs undersökningar utanför Forsmark
1987.
- 25 Carsten Peterson och Per-Olof Skoglund
(1988)
Kylvattnet från Ringhals 1974-86.

- 26 Bo Juhlin (1988)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1987.
- 27 Bo Juhlin och Stefan Tobiasson (1988)
Recipientkontroll vid Breviksnäs fiskodling 1987.
- 28 Cecilia Ambjörn (1989)
Spridning och sedimentation av tippat lermaterial utanför Helsingborgs hamnområde.
- 29 Robert Hillgren (1989)
SMHIs undersökningar utanför Forsmark 1988.
- 30 Bo Juhlin (1989)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1988.
- 31 Erland Bergstrand och Stefan Tobiasson (1989)
Samordnade kustvattenkontrollen i Östergötland 1988.
- 32 Cecilia Ambjörn (1989)
Oceanografiska förhållanden i Brofjorden i samband med kylvattenutsläpp i Trommekilen.
- 33a Cecilia Ambjörn (1990)
Oceanografiska förhållanden utanför Vendelsöfjorden i samband med kylvattenutsläpp.
- 33b Eleonor Marmefelt och Jonny Svensson (1990)
Numerical circulation models for the Skagerrak - Kattegat. Preparatory study.
- 34 Kjell Wickström (1990)
Oskarshamnsverket - kylvattenutsläpp i havet - slutrapport.
- 35 Bo Juhlin (1990)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1989.
- 36 Bertil Håkansson och Mats Moberg (1990)
Glommaälvens spridningsområde i nord-östra Skagerrak
- 37 Robert Hillgren (1990)
SMHIs undersökningar utanför Forsmark 1989.
- 38 Stig Fonselius (1990)
Skagerrak - the gateway to the North Sea.
- 39 Stig Fonselius (1990)
Skagerrak - porten mot Nordsjön.
- 40 Cecilia Ambjörn och Kjell Wickström (1990)
Spridningsundersökningar i norra Kalmarsund för Mönsterås bruk.
- 41 Cecilia Ambjörn (1990)
Strömningsteknisk utredning avseende utbyggnad av gipsdeponi i Landskrona.
- 42 Cecilia Ambjörn, Torbjörn Grafström och Jan Andersson (1990)
Spridningsberäkningar - Klints Bank.
- 43 Kjell Wickström och Robert Hillgren (1990)
Spridningsberäkningar för EKA-NOBELS fabrik i Stockviksverken.
- 44 Jan Andersson (1990)
Brofjordens kraftstation - Kylvattenspridning i Hanneviken.
- 45 Gustaf Westring och Kjell Wickström (1990)
Spridningsberäkningar för Höganäs kommun.
- 46 Robert Hillgren och Jan Andersson (1991)
SMHIs undersökningar utanför Forsmark 1990.
- 47 Gustaf Westring (1991)
Brofjordens kraftstation - Kompletterande simulering och analys av kylvattenspridning i Trommekilen.
- 48 Gustaf Westring (1991)
Vågmätningar utanför Kristianopel - Slutrapport.
- 49 Bo Juhlin (1991)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1990.
- 50A Robert Hillgren och Jan Andersson (1992)
SMHIs undersökningar utanför Forsmark 1991.

- 50B Thomas Thompson, Lars Ulander, Bertil Håkansson, Bertil Brusmark, Anders Carlström, Anders Gustavsson, Eva Cronström och Olov Fäst (1992). BEERS -92. Final edition.
- 51 Bo Juhlin (1992)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1991.
- 52 Jonny Svensson och Sture Lindahl (1992)
Numerical circulation model for the Skagerrak - Kattegat.
- 53 Cecilia Ambjörn (1992)
Isproppsförebyggande muddring och dess inverkan på strömmarna i Torneälven.
- 54 Bo Juhlin (1992)
20 års mätningar längs svenska kusten med kustbevakningens fartyg (1970 - 1990).
- 55 Jan Andersson, Robert Hillgren och Gustaf Westring (1992)
Förstudie av strömmar, tidvatten och vattenstånd mellan Cebu och Leyte, Filippinerna.
- 56 Gustaf Westring, Jan Andersson, Henrik Lindh och Robert Axelsson (1993)
Forsmark - en temperaturstudie. Slutrapport.
- 57 Robert Hillgren och Jan Andersson (1993)
SMHIs undersökningar utanför Forsmark 1992.
- 58 Bo Juhlin (1993)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1992.
- 59 Gustaf Westring (1993)
Isförhållandena i svenska farvatten under normalperioden 1961-90.
- 60 Torbjörn Lindkvist (1994)
Havsområdesregister 1993.
- 61 Jan Andersson och Robert Hillgren (1994)
SMHIs undersökningar utanför Forsmark 1993.
- 62 Bo Juhlin (1994)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1993.
- 63 Gustaf Westring (1995)
Isförhållanden utmed Sveriges kust - isstatistik från svenska farleder och farvatten under normalperioderna 1931-60 och 1961-90.
- 64 Jan Andersson och Robert Hillgren (1995)
SMHIs undersökningar utanför Forsmark 1994.
- 65 Bo Juhlin (1995)
Oceanografiska observationer runt svenska kusten med kustbevakningens fartyg 1994.
- 66 Jan Andersson och Robert Hillgren (1996)
SMHIs undersökningar utanför Forsmark 1995.
- 67 Lennart Funkquist och Patrik Ljungemyr (1997)
Validation of HIROMB during 1995-96.
- 68 Maja Brandt, Lars Edler och Lars Andersson (1998)
Översvämningar längs Oder och Wisla sommaren 1997 samt effekterna i Östersjön.
- 69 Jörgen Sahlberg SMHI och Håkan Olsson, Länsstyrelsen, Östergötland (2000).
Kustzonmodell för norra Östergötlands skärgård.
- 70 Barry Broman (2001)
En vågatlas för svenska farvatten.
Ej publicerad
- 71 ***Vakant – kommer ej att utnyttjas!***
- 72 Fourth Workshop on Baltic Sea Ice Climate Norrköping, Sweden 22-24 May, 2002
Conference Proceedings
Editors: Anders Omstedt and Lars Axell
- 73 Torbjörn Lindkvist, Daniel Björkert, Jenny Andersson, Anders Gyllander (2003)
Djupdata för havsområden 2003
- 74 Håkan Olsson, SMHI (2003)
Erik Årnefelt, Länsstyrelsen Östergötland
Kustzonssystemet i regional miljöanalys
- 75 Jonny Svensson och Eleonor Marmefelt (2003)
Utvärdering av kustzonmodellen för norra Östergötlands och norra Bohusläns skärgårdar

- 76 Eleonor Marmefelt, Håkan Olsson, Helma Lindow och Jonny Svensson, Thalassos Computations (2004)
Integrerat kustzonssystem för Bohusläns skärgård
- 77 Philip Axe, Martin Hansson och Bertil Håkansson (2004)
The national monitoring programme in the Kattegat and Skagerrak
- 78 Lars Andersson, Nils Kajrup och Björn Sjöberg (2004)
Dimensionering av det nationella marina pelagialprogrammet
- 79 Jörgen Sahlberg (2005)
Randdata från öppet hav till kustzonsmodellerna (Exemplet södra Östergötland)
- 80 Eleonor Marmefelt, Håkan Olsson (2005)
Integrerat Kustzonssystem för Hallandskusten
- 81 Tobias Strömberg (2005)
Implementation of a Flux Corrected Transport scheme in the Rossby Centre Ocean model
- 82 Martin Hansson (2006)
Cyanobakterieblomningar i Östersjön, resultat från satellitövervakning 1997-2005
- 83 Kari Eilola, Jörgen Sahlberg (2006)
Model assessment of the predicted environmental consequences for OSPAR problem areas following nutrient reductions
- 84 Torbjörn Lindkvist, Helma Lindow (2006)
Fyrskottsdata. Resultat och bearbetningsmetoder med exempel från Svenska Björn 1883 – 1892
- 85 Pia Andersson (2007)
Ballast Water Exchange areas – Prospect of designating BWE areas in the Baltic Proper
- 86 Elin Almroth, Kari Eilola, M. Skogen, H. Søiland and Ian Sehested Hansen (2007)
The year 2005. An environmental status report of the Skagerrak, Kattegat and North Sea
- 87 Eleonor Marmefelt, Jörgen Sahlberg och Marie Bergstrand (2007)
HOME Vatten i södra Östersjöns vattendistrikt. Integrerat modellsystem för vattenkvalitetsberäkningar
- 88 Pia Andersson (2007)
Ballast Water Exchange areas – Prospect of designating BWE areas in the Skagerrak and the Norwegian Trench
- 89 Anna Edman, Jörgen Sahlberg, Niclas Hjerdt, Eleonor Marmefelt och Karen Lundholm (2007)
HOME Vatten i Bottenvikens vattendistrikt. Integrerat modellsystem för vattenkvalitetsberäkningar
- 90 Niclas Hjerdt, Jörgen Sahlberg, Eleonor Marmefelt och Karen Lundholm (2007)
HOME Vatten i Bottenhavets vattendistrikt. Integrerat modellsystem för vattenkvalitetsberäkningar
- 91 Elin Almroth, Morten Skogen, Ian Sehested Hansen, Tapani Stipa, Susa Niiranen (2008)
The year 2006
An Eutrophication Status Report of the North Sea, Skagerrak, Kattegat and the Baltic Sea
A demonstration Project
- 92 Pia Andersson, editor and co-authors Bertil Håkansson*, Johan Håkansson*, Elisabeth Sahlsten*, Jonathan Havenhand**, Mike Thorndyke**, Sam Dupont** * Swedish Meteorological and Hydrological Institute ** Sven Lovén, Centre of Marine Sciences (2008)
Marine Acidification – On effects and monitoring of marine acidification in the seas surrounding Sweden
- 93 Jörgen Sahlberg, Eleonor Marmefelt, Maja Brandt, Niclas Hjerdt och Karen Lundholm (2008)
HOME Vatten i norra Östersjöns vattendistrikt. Integrerat modellsystem för vattenkvalitetsberäkningar.
- 94 David Lindstedt (2008)
Effekter av djupvattenomblandning i Östersjön – en modellstudie
- 95 Ingemar Cato*, Bertil Håkansson**, Ola Hallberg*, Bernt Kjellin*, Pia Andersson**, Cecilia Erlandsson*, Johan Nyberg*, Philip Axe** (2008)
*Geological Survey of Sweden (SGU)
**The Swedish Meteorological and Hydrological Institute (SMHI)
A new approach to state the areas of oxygen deficits in the Baltic Sea

- 96 Kari Eilola, H.E. Markus Meier, Elin Almroth, Anders Höglund (2008)
Transports and budgets of oxygen and phosphorus in the Baltic Sea
- 97 Anders Höglund, H.E. Markus Meier, Barry Broman och Ekaterini Kriezi (2009)
Validation and correction of regionalised ERA-40 wind fields over the Baltic Sea using the Rossby Centre Atmosphere model RCA3.0
- 98 Jörgen Sahlberg (2009)
The Coastal Zone Model
- 99 Kari Eilola (2009)
On the dynamics of organic nutrients, nitrogen and phosphorus in the Baltic Sea
- 100 Kristin I. M. Andreasson (SMHI), Johan Wikner (UMSC), Berndt Abrahamsson (SMF), Chris Melrose (NOAA), Svante Nyberg (SMF) (2009)
Primary production measurements – an intercalibration during a cruise in the Kattegat and the Baltic Sea
- 101 K. Eilola, B. G. Gustafson, R. Hordoir, A. Höglund, I. Kuznetsov, H.E.M. Meier T. Neumann, O. P. Savchuk (2010)
Quality assessment of state-of-the-art coupled physical-biogeochemical models in hind cast simulations 1970-2005
- 102 Pia Andersson (2010)
Drivers of Marine Acidification in the Seas Surrounding Sweden
- 103 Jörgen Sahlberg, Hanna Gustavsson (2010)
HOME Vatten i Mälaren
- 104 K.V Karmanov., B.V Chubarenko, D. Domnin, A. Hansson (2010)
Attitude to climate changes in everyday management practice at the level of Kaliningrad region municipalities
- 105 Helén C. Andersson., Patrik Wallman, Chantal Donnelly (2010)
Visualization of hydrological, physical and biogeochemical modelling of the Baltic Sea using a GeoDomeTM
- 106 Maria Bergelo (2011)
Havsvattenståndets påverkan längs Sveriges kust – enkätsvar från kommuner, räddningstjänst, länsstyrelser och hamnar
- 107 H.E. Markus Meier, Kari Eilola (2011)
Future projections of ecological patterns in the Baltic Sea
- 108 Meier, H.E.M., Andersson, H., Dieterich, C., Eilola, K., Gustafsson, B., Höglund, A., Hordoir, R., Schimanke, S (2011)
Transient scenario simulations for the Baltic Sea Region during the 21st century



Sveriges meteorologiska och hydrologiska institut
601 76 Norrköping
Tel 011-495 80 00 Fax 011-495 80 01

ISSN 0283-7714