

Report from the SMHI monitoring cruise with R/V Aranda



Survey period: 2015-10-12 - 2015-10-19

Survey area: Skagerrak, Kattegat, the Sound, the Baltic Proper and the Gulf of Finland

Principal: SMHI and the Swedish Agency for Marine and Water Management

SUMMARY

The expedition was part of the Swedish regular marine monitoring programme and covered the Skagerrak, the Kattegat, the Sound, the Baltic Proper and the Gulf of Finland. Data presented in this report has been subject to preliminary quality control procedures only.

The surface water in Skagerrak and Kattegat was somewhat cooler, around 11°C, than the surface water in the Baltic Proper where there was about 13°C. Inorganic nitrogen (NO₂+NO₃) in surface waters were close to the detection limit in all regions. The concentrations of phosphate were low throughout the region except in the Southern Baltic, where they were somewhat higher than normal. The effect of the large inflow event in December 2014 could not be seen farther north than the station BY20 in the Eastern Gotland Basin. The inflow event contributed to more oxygen in the Baltic Proper deep water but oxygen levels are now declining and the oxygen concentration in the Gotland Deep had now been reduced to 0.5 ml/l. The concentration of hydrogen sulphide has decreased which probably is an effect of the inflow. In Western- and Northern Gotland Basins the oxygen situation remains severe. In the Western Gotland Basin, completely oxygen-free conditions were found from 80-90 meters while acute hypoxia occurred from 80 meters. The northern part was completely oxygen-free from 70-80 meters and acute hypoxia occurred from 70 meters. Oxygen levels in the bottom water in the Bornholm Basin and Hanö Bight had decreased by a further 0.5 ml/l since the previous survey in September and acute hypoxia was also noted. During the entire expedition an outflow from the Baltic to the Kattegat was ongoing and there was a strong two-layer stratification in the Sound.

The next regular cruise is scheduled to start November 9, 2015.

Address:
Sven Källfelts gata 15
SE-426 71 Vastra Frolunda
SWEDEN

Telephone:
+46 11 4958000

E-mail:
karin.wesslander@smhi.se
WWW:
<http://www.smhi.se/>

PRELIMINARY RESULTS

The expedition was conducted aboard the Finnish research vessel Aranda and started in Helsinki on October 12 and ended in the same port on October 19. The winds during the expedition were mainly easterly and weak. Air temperatures ranged between 5.6 and 11.3 °C. During the entire expedition an outflow from the Baltic to the Kattegat was ongoing.

In the Gulf of Finland and the northern Baltic Proper four stations were visited also this time that usually are sampled by the Finnish Environment Institute (SYKE). This extended monitoring is part of a new collaboration between SYKE and SMHI with a view to, i.e., increasing the sampling frequency at Swedish and Finnish monitoring stations.

During the expedition a bottom-mounted measuring rig was taken up at the southern Öland. The system was deployed April 25, 2015 and has since then measured the temperature, salinity and oxygen at 58 meters depth with a measurement frequency of 10 minutes. The purpose of this operation is to follow the oxygen development in the Western Gotland Basin.

The Skagerrak

The temperature of the surface water had begun to drop and was about normal or slightly below, in the range between 11.07 and 12.29 °C. Below the cooled surface water a layer of warmer summer water was found and there below a thermocline at 40-60 meters. The salinity of the surface layer varied between 22.81 and 24.67 psu, lowest at the coast. In off-shore Skagerrak, the halocline ranged over a large depth range, 10-40 m, and surface salinity was slightly lower than normal. Closest to the coast the halocline was more developed, 10-20 m, and surface salinity around normal.

All nutrients in the surface waters showed low concentrations both at the coast and in the outer parts which is normal for the season. The concentrations in surface waters for phosphate was in the range 0.07-0.13 µmol/l, inorganic nitrogen (NO₂ + NO₃) around the detection limit <0.1 µmol/l and silicate 0.8-2.1 µmol/l. The lowest oxygen levels in the bottom water was measured at Släggö in Gullmarsfjorden estuary, 4.69 ml/l at 69 m, which is slightly higher than normal. Slightly lower oxygen concentration, 4.27 ml/l were found at the same station at 20-30 m depth is probably due the warmer water layer there.

Fluorescence measurements showed low biological activity in the surface layer 0-10 m. For more details on species composition see separate report on the algae situation.

The Kattegat and the Sound

In the Kattegat, the temperature of the surface water was in the range of 10.89 to 11.92 °C, which is normal to slightly below normal for the season. The salinity of the surface layer varied between 19.48 and 22.7 psu in the Kattegat, which is typical. In the Sound the salinity was low, 8.49 psu, depending on the outflowing surface water from the Baltic Sea. The stratification, where the halocline and thermocline coincided, were found in 10 to 20 meters depth.

The concentrations of nutrients in surface waters were normal for the season in Kattegat. The phosphate content ranged from 0.10 to 0.13 µmol/l and thus continues to rise, which is the normal case in the fall. Silicate showed lower levels than normal, 0.9-1.6 µmol/l, and inorganic nitrogen (NO₃ + NO₂) were below the reporting limit. In the Sound surface water where the phosphate concentration (0.35 µmol/l) and the content of inorganic nitrogen (0.28 µmol/l) normal and the content of silicate (5.0 µmol/l) low. However, ammonium was present (0.47 µmol/l) indicating degradation of organic material in the surface layer. In the Sound, there was also a very strong

stratification at 10 m, where the surface water had 8.49 psu in salinity and bottom water was 33.89 psu. In the stratification an oxygen-minima was observed of 2.30 ml/l, as well as elevated levels of all nutrients. The lowest oxygen levels in the bottom water were measured at Anholt E in the Kattegat (2.92 ml/l) and at W Landskrona in the Sound (2.84 ml/l).

Plankton activity was low in the surface layer 0-10 m. For details about the species composition see separate report on the algae situation.

The Baltic Proper

The temperature in the surface layer were normal or slightly above normal for the season and varied from 12.14 °C to 13.88 °C, eastern and northern parts were slightly warmer than the western part. Surface salinity was normal for the time of year with slightly higher levels in the eastern part and ranged from 5.54 psu in the central Gulf of Finland to 8.03 psu in the south western Baltic. The halocline was found at 40-70 meters depth in the western and 60-80 meters depth in the eastern Gotland Basin, while it was slightly shallower in the south. In the Bornholm Basin the halocline was found at 55-65 m depth. Thermocline was at 30 meters depth and was well developed.

The concentrations of phosphate in surface waters were now higher than in September throughout the Baltic Sea. The concentrations ranged between 0.12 µmol/l in the northern Baltic Proper to 0.49 µmol/l in the south western Baltic Proper where levels were somewhat higher than normal. There was still somewhat elevated levels of silicate which varied between 7.1 and 12.2 µmol/l. The highest phosphate levels were noted in the Arkona Basin and the highest silicate at the coastal station REFM1V1. Inorganic nitrogen (NO₂ + NO₃) was still completely exhausted in the whole area except at the coastal station RefM1V1 where it could be measured inorganic nitrogen in surface waters.

In order to follow the progress of the major inflow to the Baltic Sea, which occurred in December 2014, was also during this expedition an additional sampling carried out at station BY21 in Northern Gotland Basin. It appeared, however, no effects of the inflow event at this station. Just like the last sampling in September, an effect of the inflow at BY20 was still visible. Hydrogen sulphide was still present from 80 meters depth, but with slightly lower concentrations than during the previous survey.

In Western- and Northern Gotland Basins the oxygen situation remains very severe. In the Western Gotland Basin completely oxygen-free conditions, when hydrogen sulphide is formed, were found from 80-90 m while acute hypoxia, <2 ml/l, were found from 80 meters. The northern part was completely oxygen-free from 70-80 meters and had acute hypoxia from 70 meters.

In the Gotland Deep in the Eastern Gotland Basin, acute hypoxia occurred at depths exceeding 80 meters. The entire water column was now oxygenated, and no hydrogen sulphide was measured. Concentration of oxygen, however, had further declined slightly since the previous expedition in September and was now around 0.5 ml/l. Oxygen levels in the bottom water in the Bornholm Basin and Hanö Bight had decreased by 0.5 ml/l since the previous survey in September and acute hypoxia thus remained.

Fluorescence measurements showed low biological activity of the thermocline. For more details on species composition see separate report on the algae situation.




Figure 1. Transect showing the oxygen and salinity from the Sound to the Gulf of Finland.

PARTICIPANTS


Name		Institute
Karin Wesslander	Chief scientist	SMHI
Kristin Andreasson		SMHI
Lars Andersson		SMHI
Johan Kronsell	Lysekil-Helsinki	SMHI
Daniel Simonsson	Helsinki-Lysekil	SMHI
Sari Sipilä		SMHI

APPENDICES

- Track chart
- Table over stations, parameters and sampling depths
- Map showing bottom oxygen concentrations
- Monthly average surface water plots for selected stations
- Vertical profiles for selected stations

TRACKCHART

Country: Sweden
 Ship: R/V ARANDA
 Date: 20151012-20151019
 Series: 0606-0636



Date: 2015-10-19
Time: 16:36

Seer no	Stat code	P ro j	Station-----	Lat-----	Lon-----	Date yyyymmdd	Time hhmm utc	Bottom depth m	Secchi depth m	Wind di ve	Air temp	Air pres	WCSI elec aoad	C d	PPCPZZT Hrhhhoor Cilyooa motPBw PrP l	No de	T e	S a	P h	O x	H 2	P S	T 4	N t	N t	N t	A h	S h	L i	P u	P i	T o	O o
0606	GFXK29EXT		BY23/LL7	N5950.5	E2450.3	20151012	1200	102	5	29 6	9.8 1031	1130 x	---	13	x	x	-	x	-	x	x	x	x	x	x	x	-	x	-	-	-	-	x
0607	BPXK60EXT		LL12	N5929.0	E2254.0	20151012	1755	82		31 3	9.8 1031	9990 x	---	11	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	
0608	BPXK00EXT		LL15	N5911	E2144.81	20151012	2230	132		00 0	9.9 1031	9990 x	---	15	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	
0609	BPXK00EXT		LL17	N5902	E2104.77	20151013	0100	169		11 2	10.6 1031	9990 x	---	16	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	x
0610	BPXK27EXT		BY21	N5826.5	E2020	20151013	0530	120		09 4	10.2 1030	2830 x	---	14	x	x	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
0611	BPXK26BAS		BY20 FÅRÖDJ	N5800	E1953	20151013	0850	194	7	12 8	9.6 1029	2730 x	---	17	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	x
0612	BPXK21BAS		BY15 GOTLANDSDJ	N5720	E2003	20151013	1310	238	7	11 8	9.2 1027	2730 x	-----	19	x	x	x	x	x	x	x	x	x	x	x	x	x	x	-	-	-	-	-
0613	BPXK21EXT		BY15 GOTLANDSDJ	N5720	E2003	20151013	1440	238		11 8	9.7 1027	1430 x	-----	5	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	-
0614	BPXK13BAS		BY10	N5638	E1935	20151013	1925	144		14 8	10.1 1026	9990 x	---	15	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	-
0615	BPSE11BAS		BCS III-10	N5533.3	E1824	20151014	0250	89		10 8	9.8 1024	9990 x	---	12	x	x	-	x	-	x	x	x	x	x	x	-	x	x	-	-	-	-	-
0616	BPSB07BAS		BY5 BORNHOLMSDJ	N5515	E1559	20151014	1037	90	8	10 10	9.8 1024	2830 x	-----	12	x	x	x	-	x	x	x	x	x	x	x	x	x	x	-	-	-	-	-
0617	BPSB06BAS		BY4 CHRISTIANSÖ	N5523	E1520	20151014	1355	91		10 11	10.4 1024	2830 x	---	12	x	x	-	x	x	x	x	x	x	x	x	-	x	-	-	-	-	-	-
0618	BPSA03BAS		BY2 ARKONA	N5500	E1405	20151014	1940	44		10 13	10.6 1024	9990 x	-----	7	x	x	-	x	-	x	x	x	x	x	x	-	x	x	-	-	-	-	x
0619	BPSA02BAS		BY1	N5500	E1318	20151014	2250	46		10 11	11.3 1024	9990 x	---	8	x	x	-	x	-	x	x	x	x	x	x	-	x	-	-	-	-	-	-
0620	SOCX39BAS		W LANDSKRONA	N5552.0	E1245.0	20151015	0500	50		06 8	8.1 1024	9990 x	---	9	x	x	-	x	-	x	x	x	x	x	x	-	x	-	-	-	-	-	-
0621	KAEX29BAS		ANHOLT E	N5640.0	E1207.0	20151015	1020	63	9	09 7	9 1026	4930 x	-----	10	x	x	x	-	x	x	x	x	x	x	x	x	x	x	-	-	-	-	-
0622	KANX50BAS		N14 FALKENBERG	N5656.40	E1212.70	20151015	1245																										

Bottom water oxygen concentration (ml/l)

Country: Finland

Ship : Aranda

Date : 20151012-20151018







Series : 0606-0636





STATION P2 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >75m)



Vertical profiles P2 October







— Mean 1996-2010 St.Dev. ● 2015





STATION Å13 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >=75m)



Vertical profiles Å13 October







— Mean 1996-2010 St.Dev. ● 2015





STATION Å15 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >=125m)




Vertical profiles Å15 October

— Mean 1996-2010 St.Dev. ● 2015


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ µmol/l



DIN µmol/l









SiO₃ µmol/l





STATION Å17 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth = 300m)




Vertical profiles Å17 October

— Mean 1996-2010 St.Dev. ● 2015


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ µmol/l



DIN µmol/l









SiO₃ µmol/l





STATION FLADEN SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth > 70m)



Vertical profiles Fladen October







— Mean 1996-2010 St.Dev. ● 2015





STATION ANHOLT E SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth > 50m)



Vertical profiles Anholt E October







— Mean 1996-2010 St.Dev. ● 2015





STATION W LANDSKRONA SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles W Landskrona October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY1 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles BY1 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY2 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles BY2 October







— Mean 1996-2010 St.Dev. ● 2015





STATION HANÖBUKTEN SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth > 70m)



Vertical profiles Hanöbukten October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY4 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >80m)



Vertical profiles BY4 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY5 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >80m)



Vertical profiles BY5 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BCS III-10 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth > 80m)



Vertical profiles BCS III-10 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY10 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >125m)



Vertical profiles BY10 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY15 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >225m)



Vertical profiles BY15 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY20 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >175m)




Vertical profiles BY20 October

— Mean 1996-2010 St.Dev. ● 2015


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION BY32 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth > 175m)



Vertical profiles BY32 October







— Mean 1996-2010 St.Dev. ● 2015





STATION BY38 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (> 100m)



Vertical profiles BY38 October







— Mean 1996-2010 St.Dev. ● 2015





STATION SLÄGGÖ SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015




OXYGEN IN BOTTOM WATER (depth >50m)



Vertical profiles Släggö October

— Mean 1996-2010 St.Dev. ● 2015









STATION N14 Falkenberg SURFACE WATER

Annual Cycles



— Mean 2007-2010

..... St.Dev.

● 2015




OXYGEN IN BOTTOM WATER (depth > 25m)



Vertical profiles N14 Falkenberg October







— Mean 1996-2010 St.Dev. ● 2015





STATION REF M1V1 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2015



OXYGEN IN BOTTOM WATER (depth >15m)



Vertical profiles Ref M1V1 October

— Mean 1996-2010 St.Dev. ● 2015

