

Report from the SMHI monitoring cruise with R/V Aranda



Survey period: 2014-11-08 - 2014-11-17
Survey area: Skagerrak, Kattegat, the Sound and the Baltic Proper
Principal: SMHI and the Swedish Agency for Marine and Water Management

SUMMARY

The expedition was part of the Swedish regular marine monitoring programme and covered Skagerrak, Kattegat, the Sound and the Baltic Proper. Data presented in this report have been subject to preliminary quality control procedures only.

The water temperature in the surface layer was still above normal in all areas. Surface nutrients showed, for the season, almost normal values, except for silicate in the Baltic Proper. At the station BY1 in the Arkona Basin all nutrients showed increased values. In the Bornholm Basin and Hanö Bight acute hypoxia (< 2 ml/l) was present from depths exceeding 70 meters. In the Eastern Gotland Basin completely anoxic conditions were found from depths exceeding 125 meters (at BY20 already from 80 m) and acute hypoxia from 60 - 80 meters. In the Western Gotland Basin the oxygen situation was serious as acute hypoxia already occurred from depths exceeding 70 meters and hydrogen sulphide from 80-90 meters depth.

The next cruise is planned for December 6-14 and will cover Skagerrak, Kattegat, the Sound and the Baltic Proper.

PRELIMINARY RESULTS

The cruise, performed on board the Finnish research vessel Aranda, began in Helsinki on 8th of October and ended in Turku on the 15th. Winds during the expedition were mainly fresh, around 10m/s, varying in direction between east and south. Air temperature varied between 6 - 10°C. Due to problems with a winch no CTD measurements were carried out in the northern Kattegat or Skagerrak, which means no CTD-profiles or data on salinity can be presented from these areas. Additionally, a problem with the nutrient auto analyzer entails that phosphate- and ammonium data are missing from stations in the Skagerrak.

The Skagerrak

Temperature, down to a depth of about 100 meters, was clearly higher than normal for the season. The surface water temperatures were between 10.5 and 13.5 °C. Nutrient levels in the surface layer showed concentrations normal for the season. The concentrations of inorganic nitrogen (nitrite + nitrate) varied between 0.9 and 1.3 µmol/l, while silicate ranged from 2.1 to 2.7 µmol/l. Data on phosphate are missing from the area. The lowest oxygen level in the bottom waters were measured at the station Släggö, in the mouth of the Gullmar Fjord, where oxygen levels had dropped from 3.1 ml/l at the previous occasion in October to 2.3 ml/l.

The Kattegat and the Sound

In the Kattegat, the temperature was above normal throughout the whole water column, with surface values varying in the range of 10.5 to 11.5 °C. Thermocline and halocline coincided and were found at depths between 15 and 20 meters. The salinity of the surface water was normal, about 23 psu, in the Sound 9.5 psu.

The concentrations of phosphate and inorganic nitrogen (nitrite + nitrate) showed values typical for the season, while silicate levels were lower than normal. Phosphate varied between 0.16 and 0.19 µmol/l, inorganic nitrogen from below the detection limit (<0.10 µmol/l) to 0.6 µmol/l. In the Sound levels were generally higher, phosphate 0.38 µmol/l and nitrite + nitrate at 1.7 µmol/l. Silicate concentrations in Kattegat were lower than normal, 0.7 – 2.0 while they were normal in the Sound 8.8 µmol/l.

Fluorescence measurements showed some plankton activity in the surface layer.

The lowest oxygen levels in the bottom waters were measured at Anholt E in the southern Kattegat as well as in the Sound, 3.81 ml/l.

The Baltic Proper

Water temperatures, throughout the whole surface layer down to the thermocline at 30 to 40 meters, were well above normal in the whole area, ranging from 8.7 °C in the north to 12.7 in the south. The salinity of the surface water was normal in most of the area, between 6.5 and 8.2 psu. In the Eastern Gotland Basin surface salinity is still approximately 0.15 psu below normal, 7 psu. The halocline was found at about 60 to 70 meters depth in the Western and Eastern Gotland Basins, while in the southern parts at depths between 30 and 50 meters.

Nutrients showed almost normal levels for the season in the surface layer, phosphate concentrations ranged from 0.25 to 0.39 µmol/l, while the concentrations of inorganic nitrogen (nitrite + nitrate) varied from 0.16 to 1.26 µmol/l. Silicate showed slightly elevated levels in the northern and central parts, whereas the concentration in the southern regions were below normal. The concentrations ranged between 4.7 and 12.5 µmol/l. The station BY1 in Arkona differed completely from the pattern and had clearly elevated levels of all nutrients. Phosphate concentration was 0.68 µmol/l, silicate 12.9 µmol/l and the concentrations of nitrite + nitrates was 3.05 µmol/l.



Fluorescence measurements showed that plankton activity was low throughout the area.

The effects of an inflow through the Sound of about 25 km³ that took place in October were seen at BY2 in the Arkona Basin where the salinity in the deep water was as high as 22 psu. Also in the Bornholm Basin effects were seen at BY4 where oxygen in the bottom waters was higher than in the water above, 1.45 ml/l versus 0.65 ml/l. Acute hypoxia occurred in the Bornholm Basin and Hanö Bight from a depth of 70 meters. In the central parts of the Eastern Gotland Basin acute hypoxia occurred from 60 - 80 meters depth and hydrogen sulphide at depths exceeding 125 - 135 meters. At the station BY20, in the north, hydrogen sulphide was present already at 80 meters depth. In the Western Gotland Basin the oxygen situation was serious as acute hypoxia occurred from depths exceeding 70 meters and hydrogen sulphide from 80-90 meters depth.


PARTICIPANTS

Name		Institute
Anna-Kerstin Thell	Cruise leader	SMHI
Lars Andersson		SMHI
Sara Johansson (Lysekil – Turku)		SMHI
Mikael Krysell (Helsingfors – Lysekil)		SMHI
Vivi Månsson		SMHI
Sari Sipilä		SMHI

APPENDICES

- Track chart
- Table over stations, parameters and sampling depths
- Map showing bottom oxygen concentrations
- Monthly average surface water plots for selected stations
- Vertical profiles for selected stations

TRACKCHART
 Country: Sweden
 Ship: R/V ARANDA
 Date: 20141108-20141115
 Series: 0747-0771



Date: 2014-11-15
Time: 07:05

Ser no	Stat code	P r o j	Station-----	Lat-----	Lon-----	Date yyyymmdd	Time hhmm utc	Bottom depth m	Secchi depth m	Wind direction	Air temp C	Air pres hPa	WCSI elec t aoae d	C PPCPZPT Hrhhhoor Cilyooa tuPBw Pr p l	No de e a x h y S 4 t P N N N A S H L P T C	
0747	BPNX35BAS	BY29		N5853	E2019	20141108	2300	175		19 6	8.8	1015	9990	x --x----	16	x - x x x x x x x x - x - - - - x
0748	BPNX37BAS	BY31 LANDSORTSDJ		N5835	E1814	20141109	0720	447	10	29 7	9.4	1017	4930	x -xxxx--	23	x x x x x x x x x x - - - - x
0749	BPWX38BAS	BY32 NORRKÖPINGSDJ		N5801	E1759	20141109	1310	200	10	19 6	9	1018	2730	x --x----	17	x - x x x x x x x x - - - - x
0750	BPWX45BAS	BY38 KARLSÖDJ		N5707	E1740	20141109	2010	109		15 13	9.2	1017	9990	x --x----	14	x - x x x x x x x x - - - - x
0751	BPWK01BAS	REF M1V1		N5622.25	E1612.1	20141110	0615	18		18 8	10.6	1012	2830	x -xxxx--	5	x x x x - x x x x x x x x - - - - x
0752	BPSH05BAS	HANÖBUKTEN		N5537	E1452	20141110	1325	78	7	20 7	11.1	1009	2830	x --x----	11	x - x - x x x x x x x x - - - - x
0753	BPSA03BAS	BY2 ARKONA		N5500	E1405	20141110	1850	47		22 6	10.1	1010	9990	x -xxxx--	8	x - x - x x x x x x x x - - - - x
0754	BPSA02BAS	BY1		N5500	E1318	20141110	2210	46		23 10	10.3	1011	9990	x --x----	8	x x - x - x x x x x x x x - - - - x
0755	SOCX39BAS	W LANDSKRONA		N5552.0	E1245.0	20141111	0420	50		21 5	9.5	1011	9990	x --x----	9	x x - x - x x x x x x x x - - - - x
0756	KAXE29BAS	ANHOLT E		N5640.0	E1207.0	20141111	1000	63	9	20 9	9.6	1013	2730	x -xxxx--	10	x x x - x x x x x x x x x x - - - - x
0757	KANX50BAS	N14 FALKENBERG		N5656.40	E1212.70	20141111	1212	31	8.5	19 7	9.6	1014	2830	x -xxxx--	7	x x x x - x x x x x x x x x x - - - - x
0758	KANX25BAS	FLADEN		N5711.5	E1140	20141111	1530	85		15 6	10.8	1014	2730	- --x----	12	x x - x - x x x x x x x x - - - - x
0759	SKEX23BAS	P2		N5752	E1118	20141111	2040	92		12 6	9.6	1015	9990	- --x----	10	x x - x - x x x x x x x x - - - - x
0760	SKEX18BAS	Å17		N5816.5	E1030.8	20141112	0020	353		12 6	10.4	1015	9990	- --xx---	14	x x x x - x x x x - x x x - - - - x
0761	SKEX16BAS	Å15		N5817.7	E1051	20141112	0250	135		08 10	8.1	1015	9990	- --x----	12	x x - x - x x x x x x x x - - - - x
0762	SKEX14BAS	Å13		N5820.2	E1102	20141112	0420	100		09 10	7.6	1014	9990	- --x----	10	x x - x - x x x x x x x x - - - - x
0763	FIBG27BAS	SLÄGGÖ		N5815.5	E1126.0	20141112	0645	76	5	08 10	7.2	1015	2730	- -xxx---	9	x x - x - x x x x - x x x - - - - x
0764	KAXE29BAS	ANHOLT E		N5640.0	E1207.0	20141112	1920	50		11 12	10.6	1011	9990	x -xxxx--	9	x x x - x x x x x x x x x x - - - - x
0765	BPSB06BAS	BY4 CHRISTIANSÖ		N5523	E1520	20141113	1245	91		08 10	9.1	1015	2840	x -xx----	12	x x - x - x x x x x x x x - - - - x
0766	BPSB07BAS	BY5 BORNHOLMSDJ		N5515	E1559	20141113	1535	90		09 9	8.6	1016	9990	x --xx---	12	x x x x - x x x x x x x x x x - - - - x
0767	BPSE11BAS	BCS III-10		N5533.3	E1824	20141113	2335	91		11 8	9.1	1020	9990	x --x----	12	x x - x - x x x x x x x x - - - - x
0768	BPEX13BAS	BY10		N5638	E1935	20141114	0705	143		11 9	7.5	1024	2840	x -xxx---	15	x x - x x x x x x x x x x x x - - - - x
0769	BPEX21BAS	BY15 GOTLANDSDJ		N5720	E2003	20141114	1150	239	7	13 9	7.0	1024	2840	x -xxx---	19	- x x x x x x x x x x x x x x - - - - x
0770	BPEX21EXT	BY15 GOTLANDSDJ		N5720	E2003	20141114	1245	239		13 9	7.0	1024	2840	x -----	5	- x - x x x x x x x x x x - - - - x
0771	BPEX26BAS	BY20 FARÖDJ														


Bottom water oxygen concentration (ml/l)

Country: Finland

Ship : Aranda

Date : 20141108-20141114







Series : 0747-0771





STATION P2 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >75m)



Vertical profiles P2 November







— Mean 1996-2010 St.Dev. ● 2014





STATION Å13 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >=75m)



Vertical profiles Å13 November







— Mean 1996-2010 St.Dev. ● 2014





STATION Å15 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >=125m)




Vertical profiles Å15 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ µmol/l



DIN µmol/l









SiO₃ µmol/l





STATION Å17 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth = 300m)




Vertical profiles Å17 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION FLADEN SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth > 70m)



Vertical profiles Fladen November







— Mean 1996-2010 St.Dev. ● 2014





STATION ANHOLT E SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth > 50m)



Vertical profiles Anholt E November







— Mean 1996-2010 St.Dev. ● 2014





STATION W LANDSKRONA SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles W Landskrona November







— Mean 1996-2010 St.Dev. ● 2014





STATION BY1 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles BY1 November







— Mean 1996-2010 St.Dev. ● 2014





STATION BY2 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >40m)



Vertical profiles BY2 November







— Mean 1996-2010 St.Dev. ● 2014





STATION HANÖBUKTEN SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth > 70m)



Vertical profiles Hanöbukten November







— Mean 1996-2010 St.Dev. ● 2014





STATION BY4 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >80m)




Vertical profiles BY4 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION BY5 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >80m)



Vertical profiles BY5 November







— Mean 1996-2010 St.Dev. ● 2014





STATION BCS III-10 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth > 80m)



Vertical profiles BCS III-10 November







— Mean 1996-2010 St.Dev. ● 2014





STATION BY10 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >125m)



Vertical profiles BY10 November







— Mean 1996-2010 St.Dev. ● 2014





STATION BY15 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >225m)




Vertical profiles BY15 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION BY20 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >175m)




Vertical profiles BY20 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION BY32 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth > 175m)




Vertical profiles BY32 November

— Mean 1996-2010 St.Dev. ● 2014


Temperature °C




Salinity psu




Oxygen ml/l




PO₄ μmol/l



DIN μmol/l









SiO₃ μmol/l





STATION BY38 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (> 100m)



Vertical profiles BY38 November







— Mean 1996-2010 St.Dev. ● 2014





STATION SLÄGGÖ SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014




OXYGEN IN BOTTOM WATER (depth >50m)



Vertical profiles Släggö November

— Mean 1996-2010 St.Dev. ● 2014









STATION N14 Falkenberg SURFACE WATER

Annual Cycles



— Mean 2007-2010

..... St.Dev.

● 2014




OXYGEN IN BOTTOM WATER (depth > 25m)



Vertical profiles N14 Falkenberg November







— Mean 1996-2010 St.Dev. ● 2014





STATION REF M1V1 SURFACE WATER

Annual Cycles

— Mean 1996-2010 St.Dev. ● 2014



OXYGEN IN BOTTOM WATER (depth >15m)



Vertical profiles Ref M1V1 November

— Mean 1996-2010 St.Dev. ● 2014

