

Oceanographic Services


Lars Edler

ALGAL SITUATION IN SWEDISH MARINE WATERS

No 3, 2001, 23 April - 28 April

Quantitative samples were obtained within SMHI's regular monitoring programme, covering the Skagerrak, Kattegat, Sound and Baltic proper. The samples were scanned for toxic and dominating species of phytoplankton.

OVERVIEW



Oceanographic Services

Lars Edler

**ALGAL SITUATION IN
SWEDISH MARINE WATERS****No 3, 2001, 23 April - 28 April****DETAILS**

* POTENTIALLY HARMFUL SPECIES

SKAGERRAK**Station Å17, 23 APRIL**

Small flagellates dominated. *Monads and flagellates* less than 10 µm were most abundant. *Chrysochromulina* sp.* relatively common. Diatoms almost absent, whereas small amounts of Dinoflagellates, e.g. *Dinophysis* spp.* and *Gonyaulax grindleyi* were present.

Station Släggö, 23 APRIL

A rich flora dominated by small flagellates, including **cf. *Heterosigma*** *, with about 800 000 cells/l. Among diatoms *Thalassiosira nordenskioeldii* and *Chaetoceros constrictus* were most common. The most abundant dinoflagellates were *Peridiniella danica*, *Scrippsiella* spp. and *Gyrodinium spirale*. Single cells of *Alexandrium* cf. *tamarense* were observed.

KATTEGAT**Station Anholt E, 24 APRIL**

Poor plankton flora dominated by **cf. *Heterosigma*** *. Small flagellates, e.g. *Chrysochromulina* sp.*, *Teleaulax* sp. and *Plagioselmis* sp. were common. Dinoflagellates were dominated by *Gyrodinium spirale* with about 7 000 cells/l, *Peridiniella danica* with about 3 000 cells/l and *Lingulodinium polyedrum*. Diatoms were almost absent.

Station Anholt E, 28 APRIL

Somewhat more plankton algae compared to four days earlier. Still the dominance of **cf. *Heterosigma*** *, followed by *Chrysochromulina* sp.* and *Teleaulax* sp.. *Dinobryon balticum* was now common with about 140 000 cells/l. *Lingulodinium polyedrum* was the most abundant dinoflagellate. Except for a very low density of *Skeletonema costatum*, *Pseudo-nitzschia delicatissima* and *Thalassiosira* sp., diatoms were almost absent.

Station Anholt E, 8 MAY

Still diatoms were nearly absent. Only small numbers of *Guinardia flaccidda* were seen. Again **cf. *Heterosigma*** *, followed by *Chrysochromulina* sp.* were the most common with 85 000 and 20 000 cells/l respectively. *Lingulodinium polyedrum* was still the most abundant dinoflagellate, but other large autotrophic dinoflagellates have started to be present.

Oceanographic Services

Lars Edler

ALGAL SITUATION IN SWEDISH MARINE WATERS

No 3, 2001, 23 April - 28 April

BALTIC SEA

Arkona basin. Station BY2, 25 APRIL

Post springbloom situation with the flora turning into a flagellate community. Among dinoflagellates present were *Peridiniella catenata*, *Dinophysis acuminata**, *Amylax triacantha* and *Scrippsiella hangoei*. *Thalassiosira baltica* was the only diatom observed. Small flagellates e.g. *Pyramimonas* spp., *Plagioselmis* sp. and *Chrysochromulina* spp.* were relatively common. Single filaments of *Aphanizomenon* sp. observed.

Bornholm basin. Station BY5, 25 APRIL

The springbloom has terminated in this area and the phytoplankton were sparse. Still *Peridiniella catenata* was the most common species, accompanied by other dinoflagellates, e.g. *Protoperidinium bipes*, *Dinophysis acuminata**, *Amphidinium* sp. *Dinobryon balticum* had started to be common and single filaments of *Aphanizomenon* sp. were present. Small flagellates e.g. *Pyramimonas* spp., *Plagioselmis* sp, *Teleaulax* sp. and *Chrysochromulina* spp.* were common.

Southeast Baltic, Station BCS III 10, 25 APRIL

Also here the springbloom has terminated. Still *Peridiniella catenata* was the most common species, accompanied by several other dinoflagellates, e.g. *Protoperidinium bipes*, *Dinophysis acuminata**, *Gymnodinium* sp. *Dinobryon balticum* had started to be common and single filaments of *Aphanizomenon* sp. were present. Small flagellates e.g. *Pyramimonas* spp., *Plagioselmis* sp. and *Chrysochromulina* spp.* were common.

Eastern Gotland basin, Station BY15, 25 APRIL

A similar situation as at the previous station. However, there were still some of the springbloom diatoms present in the water, e.g. *Chaetoceros wighamii*, *Skeletonema costatum* and *Thalassiosira levanderi*.

Western Gotland basin, Station BY38, 27 APRIL

In this area the springbloom was in the late stage, but had obviously not yet terminated. *Peridiniella catenata* dominated together with *Thalassiosira baltica* and *Thalassiosira levanderi*. Small flagellates e.g. *Pyramimonas* spp., *Plagioselmis* sp and *Chrysochromulina* spp.* were common.